2024 PRIOR AUTHORIZATION CRITERIA

TABLE OF CONTENTS

Abiraterone Acetate	255
Abraxane	151
Actimmune	12
Acyclovir	13
Adcetris	151
Adempas	231
Aimovig	14
Akeega	255
Alecensa	255
Aliqopa	151
Alosetron Hydrochloride	15
Alunbrig	255
Alymsys	151
Alyq	241
Ambrisentan	233
Apokyn	27
Apomorphine Hydrochloride	27
Aranesp Albumin Free	118
Arcalyst	28
Aripiprazole	25
Aripiprazole Odt	
Armodafinil	
Asenapine Maleate Sl	25
Atovaquone	32
Augtyro	255
Auryxia	33
Avastin	151
Avita	
Avonex	176
Avonex Pen	
Ayvakit	255
Balversa	255
Bavencio	195
Belbuca	203
Beleodaq	151
Belsomra	34
Benlysta IV	35
Benlysta SC	37
Benztropine Mesylate	146
Besponsa	151
Besremi	255

Betaseron	177
Bexarotene Cap	255
Bexarotene Gel	45
Blenrep	151
Blincyto	151
Bortezomib	151
Bosentan	235
Bosulif	255
Braftovi	255
Brukinsa	255
Budesonide	84
Budesonide Er	85
Buprenorphine	203
Bydureon Bcise	86
Byetta	87
Cabometyx	255
Calquence	255
Caprelsa	
Carglumic Acid	
Cerezyme	
Chenodal	
Chlorpromazine Hydrochloride	
Chorionic Gonadotropin	
Cinacalcet Hydrochloride	
Cinryze	
Clemastine Fumarate	
Clobazam	39
Clorazepate Dipotassium	
Clozapine	
Cometrig	
Copaxone	
Copiktra	
Corlanor	98
Cosentyx	47
Cosentyx Sensoready Pen	
Cosentyx Uno Inj	47
Cotellic	
Cresemba	
Crysvita	
Cyltezo	
, Cyltezo Pen	
Cyltezo Pen-Cd/Uc/Hs Starter Pack	
Cyltezo Pen-Ps Starter Pack	
Cyproheptadine Hydrochloride	
Cyramza	
Cystadrops	
Cystagon	
~ , ~ · · · · · · · · · · · · · · · · ·	

Cystaran	103
Dalfampridine Er	105
Danazol	17
Danyelza	151
Darzalex	151
Darzalex Faspro	151
Daurismo	255
Dayvigo	106
Deferasirox (Exjade)	154
Deferasirox (Jadenu)	155
Depo-Testosterone	18
Dexmethylphenidate Hydrochloride	127
Diazepam	41
Diazepam Intensol	41
Diclofenac Sodium 3% Gel	282
Dicyclomine Hydrochloride	146
Dihydroergotamine Mesylate Spray	174
Dimethyl Fumarate	
Dimethyl Fumarate Starterpack	178
Diphenoxylate Hydrochloride/Atropine Sulfate	
Doxorubicin Hydrochloride Liposomal	
Droxidopa	
Dupixent	
Elelyso	
Eligard	
Emgality	110
Empliciti	151
Emsam	
Enbrel	51
Enbrel Mini	
Enbrel Sureclick	51
Endari	114
Enhertu	151
Epclusa	115
Epidiolex	
Frbitux	
Erivedge	255
Erleada	
Erlotinib Hydrochloride	
Everolimus	
Exkivity	
Eysuvis	
, Fanapt	
Fanapt Titration Pack	
Farydak	
Fasenra	
Fasenra Pen	
Fentanyl	

Fentanyl Citrate Oral Transmucosal	125
Fingolimod	179
Fintepla	126
Fluphenazine Decanoate	25
Fluphenazine Hydrochloride	25
Folotyn	151
Forteo	278
Fotivda	255
Fruzagla	255
Fulphila	92
Fulvestrant	
Gammagard Liquid	
Gammagard S/D Iga Less Than 1Mcg/MI	
Gammaplex	
Gamunex-C	
Gattex	
Gavreto	
Gazyva	
Gefitinib	
Gilotrif	
Glatiramer Acetate	
Glatopa	
Glydo	
Granix	
Haegarda	
Halaven	
Haloperidol	
Haloperidol Decanoate	
Haloperidol Lactate	
·	
Harvoni	
Herceptin	
Herceptin Hylecta	
Herzuma	
Humira	53
Humira Pediatric Crohns Disease Starter Pack	
Humira Pen	
Humira Pen-Cd/Uc/Hs Starter	
Humira Pen-Pediatric Uc Starter Pack	
Humira Pen-Ps/Uv Starter	
Hydrocodone Bitartrate Er	
Hydroxyzine Hydrochloride	
Hydroxyzine Pamoate	
Ibrance	
Icatibant Acetate	
Iclusig	
Idhifa	
llaris	
Imatinib Mesylate	255

Imbruvica	255
Imfinzi	196
Imiquimod	149
Inbrija	150
Inlyta	255
Ingovi	255
Inrebic	255
Ivermectin Cream	156
Ivermectin Tablet	157
lwilfin	255
Jakafi	255
Javygtor	
Jaypirca	
Jemperli	
Jevtana	
Kadcyla	
Kalydeco	
Kanjinti	
Kerendia	
Keytruda	
Kineret	
Kisqali	
Kisqali Femara 200 Dose	
Kisqali Femara 400 Dose	
Kisqali Femara 600 Dose	
Korlym	
Koselugo	
Krazati	
Kyprolis	
Lapatinib Ditosylate	
Ledipasvir/Sofosbuvir	
Lenalidomide	
Lenvima 10 Mg Daily Dose	
Lenvima 12Mg Daily Dose	256
Lenvima 14 Mg Daily Dose	
Lenvima 18 Mg Daily Dose	
Lenvima 20 Mg Daily Dose	
Lenvima 24 Mg Daily Dose	
Lenvima 4 Mg Daily Dose	
Lenvima 8 Mg Daily Dose	
Leukine	94
Leuprolide Acetate	
Libtayo	
Lidocaine Hcl Gel	162
Lidocaine Hcl Jelly	
Lidocaine Ointment	163
Lidocaine Patch	164
Lidocaine Solution	165

Lidocaine/Prilocaine	166
Lidocan	164
Lidocan III	164
Linezolid	168
Lonsurf	256
Lorazepam	42
Lorazepam Intensol	42
Lorbrena	256
Loxapine	25
Lumakras	256
Lumoxiti	151
Lupron Depot (1-Month)	161
Lupron Depot (3-Month)	161
Lupron Depot (4-Month)	161
Lupron Depot (6-Month)	
Lupron Depot-Ped (1-Month)	
Lupron Depot-Ped (3-Month)	
Lupron Depot-Ped (6-Month)	
Lurasidone	
Lybalvi	
Lynparza	
Lytgobi	
Margenza	
Matulane	
Mayzent	
Mayzent Starter Pack	
Mekinist	
Mektovi	
Memantine Hcl Titration Pak	
Memantine Hydrochloride	
Memantine Hydrochloride Er	
Methscopolamine Bromide	
Methylphenidate Hydrochloride (Methylin)	
Methylphenidate Hydrochloride (Ritalin)	250
Methylphenidate Hydrochloride Er Tablet	
Methyltestosterone	
Mifepristone	
Miglustat	
Modafinil	
Molindone Hydrochloride	
Monjuvi	
Morphine Sulfate Er	
Mvasi	
Myalept	
Mylotarg	
Natpara	
Nelarabine	
Nerlynx	
INCLIATIVE	

Ninlaro	256
Nivestym	95
Noxafil Pak	224
Nubega	256
Nuedexta	188
Nuplazid	189
Nurtec	
Ocaliva	
Octreotide Acetate	
Odomzo	
Ofev	
Ogivri	
Ogsiveo	
Ojjaara	
Olanzapine	
Olanzapine Odt	
Omnitrope	
·	
Onivyde	
Ontruzant	
Onureg	
Opdivo	
Opsumit	
Oralair	
Orencia	
Orencia Clickject	
Orgovyx	
Orkambi	
Orserdu	
Otezla	217
Oxazepam	43
Oxycontin	211
Ozempic	219
Padcev	152
Paliperidone Er	25
Palynziq	220
Panretin	221
Pazopanib	256
Pegasys	222
Pemazyre	256
Pemetrexed	
Perjeta	
Perphenazine	
Phesgo	
Pimecrolimus	
Pimozide	
Pigray 200Mg Daily Dose	
Pigray 250Mg Daily Dose	
Pigray 300Mg Daily Dose	
ı iylay Juulviş Daliy Duje	∠30

Pirfenidone	223
Plegridy	182
Plegridy Starter Pack	182
Polivy	152
Pomalyst	256
Portrazza	152
Posaconazole Dr	224
Posaconazole Inj	224
Posaconazole Susp	224
Poteligeo	152
Pregnyl	90
Procrit	119
Prolastin-C	16
Prolia	226
Promacta	229
Promethazine Hcl Plain	146
Promethazine Hydrochloride	146
Promethegan	
Pyrimethamine	
, Qinlock	
Quetiapine Fumarate	
Quetiapine Fumarate Er	
Quinine Sulfate	
Regranex	
Renflexis	
Repatha	
Repatha Pushtronex System	
Repatha Sureclick	
Retacrit	
Retevmo	
Revlimid	
Rexulti	
Rezlidhia	
Riabni	
Rinvoq	
Risperidone	
Risperidone Odt	
Rituxan	
Rituxan Hycela	
Roflumilast	
Romidepsin	
Rozlytrek	
Rubraca	
Ruxience	
Rybelsus	
Rybrevant	
Rydapt	
Sajazir	
ՀԱյԱՀՈ	

Sapropterin Dihydrochloride	254
Sarclisa	152
Scemblix	257
Scopolamine	146
Secuado	25
Signifor	262
Signifor Lar	260
Sildenafil	239
Sivextro	263
Skyrizi	71
Skyrizi Pen	
Sodium Oxybate	
Sodium Phenylbutyrate	
Sofosbuvir/Velpatasvir	
Somatuline Depot	
Somavert	
Sorafenib	
Sovaldi	
Spravato 56Mg Dose	
Spravato 84Mg Dose	
Sprycel	
Stelara	
Stivarga	
Strensiq	
Sunitinib Malate	
Sympazan	
Synribo	
Tabrecta	
Tacrolimus	
Tadalafil 20Mg Tablet	
Tafinlar	
Tagrisso	
Talzenna	
Tasigna	257
Tasimelteon	
Tazarotene	
Tazorac	283
Tazverik	257
Tecentriq	201
Tepmetko	257
Teriparatide	278
Testosterone	23
Testosterone Cypionate	18
Testosterone Enanthate	20
Testosterone Pump	23
Testosterone Topical Solution	
Tetrabenazine	
Thalomid	

Thioridazine Hcl	25
Thiothixene	25
Tibsovo	257
Tobramycin Neb	281
Tolvaptan	253
Tracleer	235
Tramadol Hcl Er	213
Trazimera	152
Trelstar Mixject	285
Tremfya	75
Tretinoin 10Mg Cap	257
Tretinoin Cream, Gel	284
Trientine Hydrochloride	286
Trifluoperazine Hcl	25
Trikafta	287
Trodelvy	152
Trulicity	
Trugap	
Truxima	
Tukysa	
Turalio	
Tymlos	
Tysabri	
Ubrelvy	
Udenyca	
Udenyca Onbody	
Unituxin	
Vanflyta	
Vectibix	
Vegzelma	
Venclexta	
Venclexta Starting Pack	
Ventavis	
Versacloz	25
Verzenio	257
Viberzi	
Victoza	
Vitrakvi	
Vizimpro	
Vonjo	
Voriconazole	
Vosevi	
Votrient	
Vpriv	
Vumerity	
Vyndamax	
Vyndagel	
Vyxeos	

Welireg	257
Xalkori	257
Xeljanz Solution	79
Xeljanz Tablet	80
Xeljanz Xr	81
Xermelo	300
Xgeva	301
Xifaxan	303
Xolair	304
Xospata	257
Xpovio	257
Xpovio 100 Mg Once Weekly	257
Xpovio 40 Mg Once Weekly	257
Xpovio 40 Mg Twice Weekly	257
Xpovio 60 Mg Once Weekly	257
Xpovio 60 Mg Twice Weekly	257
Xpovio 80 Mg Once Weekly	257
Xpovio 80 Mg Twice Weekly	257
Xtandi	257
Yargesa	275
Yervoy	202
Yondelis	152
Yonsa	257
Zaltrap	152
Zejula	
Zelboraf	
Zepatier	307
Zepzelca	152
Ziextenzo	97
Ziprasidone Mesylate	25
Zirabev	
Zokinvy	308
Zolinza	
Ztlido	167
Zydelig	
Zykadia	
Zynlonta	
Zynrexa Relnreyy	25

Actimmune PA

Drug Name(s)

Actimmune

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Acyclovir Topical PA

Drug Name(s)

Acyclovir

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Aimovig PA

Drug Name(s)

Aimovig

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of migraine AND
- 2. The requested agent is being used for migraine prophylaxis AND
- 3. Patient has 4 migraine headaches or more per month AND
- 4. ONE of the following:
 - A. Patient has tried and had an inadequate response to a conventional migraine prophylaxis agent [e.g., beta blockers (propranolol), anticonvulsants (divalproex, topiramate)] OR
 - B. Patient has an intolerance, or hypersensitivity to a conventional migraine prophylaxis agent OR
 - C. Patient has an FDA labeled contraindication to a conventional migraine prophylaxis agent AND
- 5. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of migraine AND
- 3. The requested agent is being used for migraine prophylaxis AND
- 4. Patient has had clinical benefit with the requested agent AND
- 5. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Alosetron PA

Drug Name(s)

Alosetron Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of irritable bowel syndrome with severe diarrhea (IBS-D) AND
- 2. Patient's sex is female AND
- 3. Patient exhibits at least ONE of the following:
 - a. Frequent and severe abdominal pain/discomfort OR
 - b. Frequent bowel urgency or fecal incontinence OR
 - c. Disability or restriction of daily activities due to IBS AND
- 4. Prescriber has ruled out anatomic or biochemical abnormalities of the gastrointestinal tract

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Alpha-1-Proteinase Inhibitor PA - Prolastin-C

Drug Name(s)

Prolastin-C

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of alpha-1 antitrypsin deficiency (AATD) with clinically evident emphysema AND
- 2. Patient has a pre-treatment serum alpha-1 antitrypsin (AAT) level less than 11 micromol/L (80 mg/dL by immunodiffusion or 57 mg/dL using nephelometry) AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of alpha-1 antitrypsin deficiency (AATD) with clinically evident emphysema AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Anabolic Steroid PA - Danazol

Drug Name(s)

Danazol

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Patient has an FDA labeled indication for the requested agent OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR
 - B. Prescriber has provided information in support of therapy with more than one agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Androgen Injectable PA - testosterone cypionate

Drug Name(s)

Depo-Testosterone

Testosterone Cypionate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Patient's sex is male with AIDS/HIV-associated wasting syndrome AND BOTH of the following:
 - i. ONE of the following:
 - a. Unexplained involuntary weight loss (greater than 10% baseline body weight within 12 months, or 7.5% within 6 months) OR
 - b. Body mass index less than 20 kg/m2 OR
 - c. At least 5% total body cell mass (BCM) loss within 6 months OR
 - d. BCM less than 35% of total body weight and BMI less than 27 kg/m2 AND
 - ii. All other causes of weight loss have been ruled out OR
 - B. Patient's sex is female with metastatic/inoperable breast cancer OR
 - C. Patient's sex is male with primary or secondary (hypogonadotropic) hypogonadism OR
 - D. Patient's sex is male and is an adolescent with delayed puberty AND
- 2. If the patient's sex is a male, ONE of the following:
 - A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:
 - i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR
 - ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR
 - B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:
 - i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR
 - ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND
- 3. ONE of the following:
 - A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR
 - B. Prescriber has provided information in support of therapy with more than one agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for delayed puberty, 12 months for all other indications Other Criteria:

Androgen Injectable PA - testosterone enanthate

Drug Name(s)

Testosterone Enanthate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Patient's sex is male with AIDS/HIV-associated wasting syndrome AND BOTH of the following:
 - i. ONE of the following:
 - a. Unexplained involuntary weight loss (greater than 10% baseline body weight within 12 months, or 7.5% within 6 months) OR
 - b. Body mass index less than 20 kg/m2 OR
 - c. At least 5% total body cell mass (BCM) loss within 6 months OR
 - d. BCM less than 35% of total body weight and BMI less than 27 kg/m2 AND
 - ii. All other causes of weight loss have been ruled out OR
 - B. Patient's sex is female with metastatic/inoperable breast cancer OR
 - C. Patient's sex is male with primary or secondary (hypogonadotropic) hypogonadism OR
 - D. Patient's sex is male and is an adolescent with delayed puberty AND
- 2. If the patient's sex is a male, ONE of the following:
 - A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:
 - i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR
 - ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR
 - B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:
 - i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR
 - ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND
- 3. ONE of the following:
 - A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR
 - B. Prescriber has provided information in support of therapy with more than one agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for delayed puberty, 12 months for all other indications

Androgen Oral PA

Drug Name(s)

Methyltestosterone

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Patient's sex is male with cryptorchidism OR
 - B. Patient's sex is male with hypogonadism OR
 - C. Patient's sex is male and is an adolescent with delayed puberty OR
 - D. Patient's sex is female with metastatic/inoperable breast cancer AND
- 2. If the patient's sex is male, ONE of the following:
 - A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:
 - i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR
 - ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR
 - B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:
 - i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR
 - ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND
- 3. ONE of the following:
 - A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR
 - B. Prescriber has provided information in support of therapy with more than one agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for delayed puberty, 12 months for all other indications

Androgen Topical PA

Drug Name(s)

Testosterone

Testosterone Pump

Testosterone Topical Solution

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Patient has AIDS/HIV-associated wasting syndrome AND BOTH of the following:
 - i. ONE of the following:
 - a. Unexplained involuntary weight loss (greater than 10% baseline body weight within 12 months, or 7.5% within 6 months) OR
 - b. Body mass index less than 20 kg/m2 OR
 - c. At least 5% total body cell mass (BCM) loss within 6 months OR
 - d. In men: BCM less than 35% of total body weight and BMI less than 27 kg/m2 OR
 - e. In women: BCM less than 23% of total body weight and BMI less than 27 kg/m2 AND
 - ii. All other causes of weight loss have been ruled out OR
- B. Patient's sex is male with primary or secondary (hypogonadotropic) hypogonadism AND 2. If the patient's sex is male, ONE of the following:
 - A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:
 - i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR
 - ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR
 - B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:
 - i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR
 - ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND
- 3. ONE of the following:
 - A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR
 - B. Prescriber has submitted information in support of therapy with more than one agent

Age Restriction:

Prescriber Restrictions:
Coverage Duration:
Approval will be for 12 months
Other Criteria:

Antipsychotics PA

Drug Name(s)

Aripiprazole

Aripiprazole Odt

Asenapine Maleate SI

Chlorpromazine Hydrochloride

Clozapine

Clozapine Odt

Fanapt

Fanapt Titration Pack

Fluphenazine Decanoate

Fluphenazine Hydrochloride

Haloperidol

Haloperidol Decanoate

Haloperidol Lactate

Loxapine

Lurasidone

Lybalvi

Molindone Hydrochloride

Olanzapine

Olanzapine Odt

Paliperidone Er

Perphenazine

Pimozide

Quetiapine Fumarate

Quetiapine Fumarate Er

Rexulti

Risperidone

Risperidone Odt

Secuado

Thioridazine Hcl

Thiothixene

Trifluoperazine Hcl

Versacloz

Ziprasidone Mesylate

Zyprexa Relprevy

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

PA does NOT apply to patients less than 65 years of age.

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent OR
 - c. ONE of the following:
 - i. Patient has a diagnosis other than dementia-related psychosis or dementia related behavioral symptoms OR
 - ii. Patient has dementia-related psychosis or dementia related behavioral symptoms AND BOTH of the following:
 - 1. Dementia-related psychosis is determined to be severe or the associated behavior puts the patient or others in danger AND
 - 2. Prescriber has documented that s/he has discussed the risk of increased mortality with the patient and/or the patient's surrogate decision maker

Approval authorizations will apply to the requested medication only.

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Apomorphine Inj PA

Drug Name(s)

Apokyn

Apomorphine Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. The requested agent will be used to treat acute, intermittent hypomobility, "off" episodes ("end of dose wearing off" and unpredictable "on/off" episodes) associated with advanced Parkinson's disease AND
- 2. The requested agent will be used in combination with agents used for therapy in Parkinson's disease (e.g., levodopa, dopamine agonist, monoamine oxidase B inhibitor) AND
- 3. Patient will NOT be using the requested agent in combination with a 5-HT3 antagonist (e.g., ondansetron, granisetron, dolasetron, palonosetron, alosetron)

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Arcalyst PA

Drug Name(s)

Arcalyst

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. ONE of the following:
 - A. Patient has been diagnosed with Cryopyrin-Associated Periodic Syndromes (CAPS) including Familial Cold Auto-inflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS) OR
 - B. BOTH of the following:
 - i. Patient has a diagnosis of deficiency of interleukin-1 receptor antagonist AND
 - ii. The requested agent is being used for maintenance of remission OR
 - C. BOTH of the following:
 - i. Patient has a diagnosis of recurrent pericarditis AND
 - ii. The requested agent is being used to reduce the risk of recurrence AND
- 2. Patient will NOT be using the requested agent in combination with another biologic agent

Age Restriction:

For diagnosis of CAPS including FCAS or MWS, patient is 12 years of age or over

For diagnosis of recurrent pericarditis and reduction in risk of recurrence, patient is 12 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Armodafinil PA

Drug Name(s)

Armodafinil

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another target agent (i.e., modafinil)

Age Restriction:

Patient is 17 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Atopic Dermatitis PA – Pimecrolimus

Drug Name(s)

Pimecrolimus

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ONE of the following:

- 1. Patient has a diagnosis of atopic dermatitis or vulvar lichen sclerosus AND ONE of the following:
 - A. Patient has tried and had an inadequate response to a topical corticosteroid or topical corticosteroid combination preparation (e.g., hydrocortisone, triamcinolone) OR
 - B. Patient has an intolerance or hypersensitivity to a topical corticosteroid or topical corticosteroid combination preparation OR
 - C. Patient has an FDA labeled contraindication to a topical corticosteroid or topical corticosteroid combination preparation OR
- 2. Patient has a diagnosis of facial seborrheic dermatitis associated with HIV infection AND BOTH of the following:
 - A. Patient is currently on an antiretroviral treatment regimen AND
 - B. ONE of the following:
 - i. Patient has tried and had an inadequate response to a topical corticosteroid or topical antifungal treatment (e.g., hydrocortisone, triamcinolone, ketoconazole) OR
 - ii. Patient has an intolerance or hypersensitivity to a topical corticosteroid or topical antifungal treatment OR
 - iii. Patient has an FDA labeled contraindication to a topical corticosteroid or topical antifungal treatment OR
- 3. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Atopic Dermatitis PA – Tacrolimus

Drug Name(s)

Tacrolimus

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ONE of the following:

- 1. Patient has a diagnosis of atopic dermatitis AND ONE of the following:
 - A. Patient has tried and had an inadequate response to a topical corticosteroid or topical corticosteroid combination preparation (e.g., hydrocortisone, triamcinolone) OR
 - B. Patient has an intolerance or hypersensitivity to a topical corticosteroid or topical corticosteroid combination preparation OR
 - C. Patient has an FDA labeled contraindication to a topical corticosteroid or topical corticosteroid combination preparation OR
- 2. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Atovaquone PA

Drug Name(s)

Atovaquone

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - 1. Patient has a diagnosis of mild-to-moderate Pneumocystis jirovecii pneumonia OR
 - 2. Patient is using the requested agent for prevention of Pneumocystis jirovecii pneumonia AND
 - ii. ONE of the following:
 - 1. Patient has an intolerance or hypersensitivity to trimethoprim/sulfamethoxazole (TMP/SMX) OR
 - 2. Patient has an FDA labeled contraindication to trimethoprim/sulfamethoxazole (TMP/SMX) OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Auryxia PA

Drug Name(s)

Auryxia

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used as iron replacement therapy to treat iron deficiency anemia AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

- 1. Patient has a diagnosis of hyperphosphatemia AND BOTH of the following:
 - A. Patient has chronic kidney disease AND
 - B. Patient is on dialysis

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Belsomra PA

Drug Name(s)

Belsomra

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benlysta IV PA

Drug Name(s)

Benlysta IV

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. ONE of the following:
 - a. Patient has a diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:
 - i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:
 - i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another biologic agent AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - a. Patient has diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:
 - i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:
 - i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another biologic agent AND
- 5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

For diagnosis of active systemic lupus erythematosus (SLE) disease, patient is 5 years of age or over. For diagnosis of active lupus nephritis (LN), patient is 5 years of age or over.

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benlysta SC PA

Drug Name(s)

Benlysta SC

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. ONE of the following:
 - a. Patient has a diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:
 - i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:
 - i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another biologic agent AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - a. Patient has diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:
 - i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:
 - i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another biologic agent AND
- 5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

For diagnosis of active systemic lupus erythematosus (SLE) disease, patient is 18 years of age or over. For diagnosis of active lupus nephritis (LN), patient is 18 years of age or over.

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA - Clobazam

Drug Name(s)

Clobazam

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. BOTH of the following:
 - i. Patient has ONE of the following diagnoses:
 - a. Seizure disorder OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Clorazepate

Drug Name(s)

Clorazepate Dipotassium

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. BOTH of the following:
 - i. Patient has ONE of the following diagnoses:
 - a. Seizure disorder OR
 - b. Anxiety disorder AND ONE of the following:
 - 1) Patient has tried and has an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR
 - 2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR
 - 3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR
 - c. Alcohol withdrawal OR
 - d. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA - Diazepam

Drug Name(s)

Diazepam

Diazepam Intensol

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. BOTH of the following:
 - i. Patient has ONE of the following diagnoses:
 - a. Seizure disorder OR
 - b. Anxiety disorder AND ONE of the following:
 - 1) Patient has tried and had an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR
 - 2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR
 - 3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR
 - c. Skeletal muscle spasms OR
 - d. Alcohol withdrawal OR
 - e. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Lorazepam

Drug Name(s)

Lorazepam

Lorazepam Intensol

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. BOTH of the following:
 - i. Patient has ONE of the following diagnoses:
 - a. Anxiety disorder AND ONE of the following:
 - 1) Patient has tried and had an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR
 - 2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR
 - 3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Oxazepam

Drug Name(s)

Oxazepam

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. BOTH of the following:
 - i. Patient has ONE of the following diagnoses:
 - a. Anxiety disorder AND ONE of the following:
 - 1) Patient has tried and had an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR
 - 2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR
 - 3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR
 - b. Alcohol withdrawal OR
 - c. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Sympazan

Drug Name(s)

Sympazan

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. BOTH of the following:
 - i. Patient has ONE of the following diagnoses:
 - a. Seizure disorder OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Bexarotene Gel PA

Drug Name(s)

Bexarotene Gel

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of stage IA or IB cutaneous T-cell lymphoma (CTCL) with cutaneous lesions OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:
 - i. ONE of the following:
 - 1. BOTH of the following:
 - a. Patient has a diagnosis of stage IA or IB cutaneous T-cell lymphoma (CTCL) with cutaneous lesions AND
 - b. ONE of the following:
 - i. Patient has refractory or persistent disease despite a previous treatment trial with a skin-directed therapy (e.g., topical corticosteroid, topical imiquimod) OR
 - ii. Patient has an intolerance or hypersensitivity to a previous treatment trial with a skin-directed therapy (e.g., topical corticosteroid, topical imiquimod) OR
 - iii. Patient has an FDA labeled contraindication to a previous treatment trial with a skin-directed therapy (e.g., topical corticosteroid, topical imiquimod) OR
 - 2. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., dermatologist, oncologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of stage IA or IB cutaneous T-cell lymphoma (CTCL) with cutaneous lesions OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Patient has had clinical benefit with the requested agent AND
 - ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., dermatologist, oncologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Biologic Immunomodulators PA - Cosentyx

Drug Name(s)

Cosentyx

Cosentyx Sensoready Pen

Cosentyx Uno Inj

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis or plaque psoriasis

NO prerequisites are required for diagnoses of ankylosing spondylitis, enthesitis related arthritis, hidradenitis suppurativa, or non-radiographic axial spondyloarthritis

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Biologic Immunomodulators PA - Cyltezo

Drug Name(s)

Cyltezo

Cyltezo Pen

Cyltezo Pen-Cd/Uc/Hs Starter Pack

Cyltezo Pen-Ps Starter Pack

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
- C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
- D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
- E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
- F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
- G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 12 weeks for initial use for ulcerative colitis, 12 months for all others **Other Criteria:**

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Crohn's disease, or moderate ulcerative colitis

NO prerequisites are required for diagnoses of ankylosing spondylitis, hidradenitis suppurativa, severe ulcerative colitis, or uveitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Biologic Immunomodulators PA - Enbrel

Drug Name(s)

Enbrel

Enbrel Mini

Enbrel Sureclick

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile psoriatic arthritis, or juvenile idiopathic arthritis

NO prerequisites are required for a diagnoses of ankylosing spondylitis or severe juvenile psoriatic arthritis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, juvenile psoriatic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Biologic Immunomodulators PA - Humira

Drug Name(s)

Humira

Humira Pediatric Crohns Disease Starter Pack

Humira Pen

Humira Pen-Cd/Uc/Hs Starter

Humira Pen-Pediatric Uc Starter Pack

Humira Pen-Ps/Uv Starter

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 12 weeks for initial use for ulcerative colitis, 12 months for all others **Other Criteria**:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Crohn's disease, or moderate ulcerative colitis

NO prerequisites are required for diagnoses of ankylosing spondylitis, hidradenitis suppurativa, severe ulcerative colitis, or uveitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Biologic Immunomodulators PA - Kineret

Drug Name(s)

Kineret

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:
 - i. Patient's diagnosis is indicated for preferred biologic immunomodulator agent(s) AND ONE of the following:
 - a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 - b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR
 - c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) $\mbox{\rm OR}$
 - ii. The request is for an FDA labeled indication that is not covered by preferred biologic immunomodulator agent(s) AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Use of TWO preferred agents (Cyltezo, Enbrel, Humira, Rinvoq, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

NO preferred agent is required for diagnoses of Neonatal-Onset Multisystem Inflammatory Disease (NOMID) or Deficiency of Interleukin-1 Receptor Antagonist (DIRA)

Biologic Immunomodulators PA - Orencia

Drug Name(s)

Orencia

Orencia Clickject

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:
 - i. Patient's diagnosis is indicated for preferred biologic immunomodulator agent(s) AND ONE of the following:
 - a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 - b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR
 - c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR
 - ii. The request is for an FDA labeled indication that is not covered by preferred biologic immunomodulator agent(s) AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of TWO preferred agents (Cyltezo, Enbrel, Humira, Xeljanz tablets, or Xeljanz oral solution) is required for diagnosis of juvenile idiopathic arthritis

Use of TWO preferred agents (Cyltezo, Enbrel, Humira, Rinvoq, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

Use of TWO preferred agents (Cosentyx, Cyltezo, Enbrel, Humira, Otezla, Rinvoq, Skyrizi, Stelara, Tremfya, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of psoriatic arthritis

NO preferred agent is required for diagnosis of prophylaxis of acute graft vs host disease

Biologic Immunomodulators PA – Renflexis

Drug Name(s)

Renflexis

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:
 - i. Patient's diagnosis is indicated for preferred biologic immunomodulator agent(s) AND ONE of the following:
 - a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 - b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR
 - c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR
 - ii. The request is for an FDA labeled indication that is not covered by preferred biologic immunomodulator agent(s) AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Use of TWO preferred agents (Cyltezo, Enbrel, or Humira) is required for diagnosis of rheumatoid arthritis

Use of TWO preferred agents (Cosentyx, Cyltezo, Enbrel, Humira, Otezla, Skyrizi, Stelara, or Tremfya) is required for diagnosis of psoriatic arthritis

Use of TWO preferred agents (Cosentyx, Cyltezo, Enbrel, Humira, Otezla, Skyrizi, Stelara, or Tremfya) is required for diagnosis of plaque psoriasis

Use of TWO preferred agents (Cosentyx, Cyltezo, Enbrel, or Humira) is required for diagnosis of ankylosing spondylitis

Use of TWO preferred agents (Cyltezo, Humira, Skyrizi, or Stelara) is required for diagnosis of adult Crohn's disease

Use of TWO preferred agents (Cyltezo, Humira, or Stelara) is required for diagnosis of adult ulcerative colitis

Use of ONE preferred agent (Cyltezo or Humira) is required for diagnosis of pediatric Crohn's disease

Only the preferred agent Humira is required for diagnosis of pediatric ulcerative colitis

NO preferred agent is required for diagnosis of adult fistulizing Crohn's disease

Biologic Immunomodulators PA - Riabni

Drug Name(s)

Riabni

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
- 2. ALL of the following:
 - A. ONE of the following:
 - i. Patient has a diagnosis of rheumatoid arthritis AND ONE of the following:
 - a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 - b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR
 - c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR
 - ii. Patient has another FDA labeled indication or an indication that is supported in CMS approved compendia AND
 - B. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND
 - C. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
 - D. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:
 - i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
 - ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
 - iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
 - iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Use of TWO preferred agents (Cyltezo, Enbrel, Humira, Rinvoq, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

ALL other diagnoses do NOT require any preferred agents

Biologic Immunomodulators PA - Rinvoq

Drug Name(s)

Rinvoq

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:
 - i. BOTH of the following:
 - a. Patient has an FDA labeled indication other than moderate to severe atopic dermatitis for the requested agent AND
 - b. ONE of the following:
 - 1. Patient's medication history indicates use of preferred TNF agent(s) OR
 - 2. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
 - 3. Patient has an FDA labeled contraindication to preferred TNF agent(s) OR
 - 4. The request is for an FDA labeled indication that is not covered by preferred TNF agent(s) OR
 - ii. Patient has a diagnosis of moderate to severe atopic dermatitis AND ONE of the following:
 - 1. Patient's medication history indicates use of TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) for the requested indication OR
 - 2. Patient has an intolerance or hypersensitivity to TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) for the requested indication OR
 - 3. Patient has an FDA labeled contraindication to TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) for the requested indication AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Use of ONE preferred TNF (Cyltezo, Enbrel, or Humira) is required for diagnoses of ankylosing spondylitis, rheumatoid arthritis, or psoriatic arthritis

Use of ONE preferred TNF (Cyltezo or Humira) is required for diagnoses of ulcerative colitis or Crohn's disease

Use of TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) are required for diagnosis of moderate to severe atopic dermatitis

NO preferred TNF agents are required for diagnosis of non-radiographic Axial Spondyloarthritis

Biologic Immunomodulators PA – Rituxan

Drug Name(s)

Rituxan

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
- 2. ALL of the following:
 - A. ONE of the following:
 - i. Patient has a diagnosis of rheumatoid arthritis AND ONE of the following:
 - a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 - b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR
 - c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR
 - ii. Patient has another FDA labeled indication or an indication that is supported in CMS approved compendia AND
 - B. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND
 - C. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
 - D. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:
 - i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
 - ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
 - iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
 - iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Use of TWO preferred agents (Cyltezo, Enbrel, Humira, Rinvoq, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

ALL other diagnoses do NOT require any preferred agents

Biologic Immunomodulators PA - Rituxan Hycela

Drug Name(s)

Rituxan Hycela

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND
 - ii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
 - iii. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following
 - i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
 - ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

- iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
- iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

There are no preferred agents required for Rituxan Hycela

Biologic Immunomodulators PA - Ruxience

Drug Name(s)

Ruxience

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
- 2. ALL of the following:
 - A. ONE of the following:
 - i. Patient has a diagnosis of rheumatoid arthritis AND ONE of the following:
 - a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 - b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR
 - c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR
 - ii. Patient has another FDA labeled indication or an indication that is supported in CMS approved compendia AND
 - B. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND
 - C. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
 - D. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:
 - i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
 - ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
 - iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
 - iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Use of TWO preferred agents (Cyltezo, Enbrel, Humira, Rinvoq, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

ALL other diagnoses do NOT require any preferred agents

Biologic Immunomodulators PA - Skyrizi

Drug Name(s)

Skyrizi

Skyrizi Pen

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - E. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of Crohn's disease, plaque psoriasis, or psoriatic arthritis

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Biologic Immunomodulators PA – Stelara

Drug Name(s)

Stelara

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, moderate ulcerative colitis, or Crohn's disease

NO prerequisites are required for diagnosis of severe ulcerative colitis

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, mercaptopurine

Biologic Immunomodulators PA – Tremfya

Drug Name(s)

Tremfya

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - E. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis or plaque psoriasis

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Biologic Immunomodulators PA - Truxima

Drug Name(s)

Truxima

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
- 2. ALL of the following:
 - A. ONE of the following:
 - i. Patient has a diagnosis of rheumatoid arthritis AND ONE of the following:
 - a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 - b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR
 - c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR
 - ii. Patient has another FDA labeled indication or an indication that is supported in CMS approved compendia AND
 - B. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND
 - C. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
 - D. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:
 - i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
 - ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
 - iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND
 - iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Use of TWO preferred agents (Cyltezo, Enbrel, Humira, Rinvoq, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

ALL other diagnoses do NOT require any preferred agents

Biologic Immunomodulators PA - Xeljanz Solution

Drug Name(s)

Xeljanz Solution

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. Patient's medication history indicates use of preferred TNF agent(s) OR
 - ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
 - iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, or Humira) is required for diagnosis of juvenile idiopathic arthritis

Biologic Immunomodulators PA - Xeljanz Tablet

Drug Name(s)

Xeljanz Tablet

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. Patient's medication history indicates use of preferred TNF agent(s) OR
 - ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
 - iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, or Humira) is required for diagnoses of psoriatic arthritis, rheumatoid arthritis, juvenile idiopathic arthritis, or ankylosing spondylitis

Use of ONE preferred TNF (Cyltezo or Humira) is required for diagnosis of ulcerative colitis

Biologic Immunomodulators PA - Xeljanz XR

Drug Name(s)

Xeljanz Xr

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. Patient's medication history indicates use of preferred TNF agent(s) OR
 - ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
 - iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) AND
- 3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
- 4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, or Humira) is required for diagnoses of psoriatic arthritis, rheumatoid arthritis, or ankylosing spondylitis

Use of ONE preferred TNF (Cyltezo or Humira) is required for diagnosis of ulcerative colitis

Bivigam/Flebogamma/Gammaplex/Octagam/Privigen PA

Drug Name(s)

Gammaplex

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ONE of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Primary immunodeficiency [e.g., congenital agammaglobulinemia, common variable immunodeficiency (CVID), severe combined immunodeficiency, Wiskott-Aldrich Syndrome, X-linked agammaglobulinemia (XLA), humoral immunodeficiency, IgG subclass deficiency with or without IgA deficiency] OR
 - B. B-cell chronic lymphocytic leukemia OR multiple myeloma AND ONE of the following:
 - i. Patient has a history of infections OR
 - ii. Patient has evidence of specific antibody deficiency OR
 - iii. Patient has hypogammaglobulinemia OR
 - C. Idiopathic thrombocytopenia purpura AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone), or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - D. Dermatomyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - E. Polymyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - F. Severe rheumatoid arthritis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., tumor necrosis factor antagonists (e.g., Humira), DMARDS (e.g., methotrexate)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions: Coverage Duration:

Approval will be for 6 months for indications in Other Criteria, 12 months for all others **Other Criteria**:

- G. Myasthenia gravis (MG) AND ONE of the following:
 - i. Patient is in acute myasthenic crisis OR
 - ii. Patient has severe refractory MG (e.g., major functional disability/weakness) AND ONE of the following:
 - a) Patient has failed ONE immunomodulator therapy (i.e., corticosteroid, pyridostigmine, or azathioprine) OR
 - b) Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE immunomodulator therapy OR
- H. Multiple sclerosis (MS) AND BOTH of the following:
 - i. Patient has a diagnosis of relapsing remitting MS (RRMS) AND
 - ii. Patient has had an insufficient response, documented failure, or FDA labeled contraindication to TWO MS agents (e.g., Avonex, Betaseron, Copaxone, dimethyl fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR
- I. Acquired von Willebrand hemophilia AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g., desmopressin solution, von Willebrand factor replacement therapy, corticosteroids, cyclophosphamide, FEIBA, or recombinant factor VIIa) OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- J. Refractory pemphigus vulgaris AND ONE of the following:
 - i. Patient has failed ONE conventional immunosuppressive therapy (e.g., azathioprine, cyclophosphamide, mycophenolate, corticosteroids) OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional immunosuppressive therapy OR
- 2. ONE of the following:
 - A. Patient has another FDA labeled indication for the requested agent OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Indications with 6 months approval duration: Acquired von Willebrand hemophilia, Guillain-Barre Syndrome, Lambert-Eaton myasthenia syndrome, Kawasaki disease, CMV induced pneumonitis in solid organ transplant, Toxic shock syndrome due to invasive group A streptococcus, Toxic epidermal necrolysis and Stevens-Johnson syndrome

Drug is also subject to Part B versus Part D review.

Budesonide Oral ER PA - Entocort

Drug Name(s)

Budesonide

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Budesonide Oral ER PA – Uceris

Drug Name(s)

Budesonide Er

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Bydureon PA

Drug Name(s)

Bydureon Bcise

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of type 2 diabetes mellitus AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 - 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Byetta PA

Drug Name(s)

Byetta

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of type 2 diabetes mellitus AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 - 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Carglumic PA

Drug Name(s)

Carglumic Acid

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA)

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of ONE of the following:
 - a. Acute hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS) OR
 - b. Chronic hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS) OR
 - c. Acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA) AND
- 2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., geneticist, nephrologist, metabolic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Chenodal PA

Drug Name(s)

Chenodal

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of radiolucent stones in a well-opacifying gallbladder AND
- 2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Chorionic Gonadotropin PA

Drug Name(s)

Chorionic Gonadotropin

Pregnyl

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used to promote fertility AND requested agent will be used to treat erectile dysfunction AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of prepubertal cryptorchidism not due to anatomic obstruction OR
 - B. Patient's sex is male, with a diagnosis of hypogonadotropic hypogonadism (hypogonadism secondary to pituitary deficiency) AND BOTH of the following:
 - i. Patient has a measured current or pretreatment total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR a free serum testosterone level that is below the testing laboratory's lower limit of the normal range AND
 - ii. Patient has measured luteinizing hormone (LH) AND follicle-stimulating hormone (FSH) levels that are at (low-normal) or below the testing laboratory's normal range OR
 - C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Cinacalcet PA

Drug Name(s)

Cinacalcet Hydrochloride

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

- 1. Patient has ONE of the following:
 - A. A diagnosis of hypercalcemia due to parathyroid carcinoma OR
 - B. A diagnosis of primary hyperparathyroidism (HPT) AND BOTH of the following:
 - i. Patient has a pretreatment serum calcium level that is above the testing laboratory's upper limit of normal AND
 - ii. Patient is unable to undergo parathyroidectomy OR
 - C. Another indication that is FDA approved or supported in CMS approved compendia for the requested agent not otherwise excluded from Part D [i.e., secondary hyperparathyroidism due to end-stage renal disease (ESRD) on dialysis]

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Colony Stimulating Factors PA – Fulphila

Drug Name(s)

Fulphila

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Granix

Drug Name(s)

Granix

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Leukine

Drug Name(s)

Leukine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Nivestym

Drug Name(s)

Nivestym

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Udenyca

Drug Name(s)

Udenyca

Udenyca Onbody

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Ziextenzo

Drug Name(s)

Ziextenzo

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Corlanor PA

Drug Name(s)

Corlanor

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has stable, symptomatic chronic heart failure (e.g., NYHA Class II, III, IV: ACCF/AHA Class C, D) AND
- 2. ONE of following:
 - a. ALL of the following:
 - i. The requested agent is for a pediatric patient, 6 months of age or over AND
 - ii. Patient has heart failure due to dilated cardiomyopathy (DCM) AND
 - iii. Patient is in sinus rhythm with an elevated heart rate OR
 - b. ALL of the following:
 - i. The requested agent is for an adult patient AND
 - ii. Patient has a baseline OR current left ventricular ejection fraction of 35% or less AND
 - iii. Patient is in sinus rhythm with a resting heart rate of 70 beats or greater per minute prior to initiating therapy with the requested agent AND
 - iv. ONE of the following:
 - 1. Patient is on a maximally tolerated dose of beta blocker (e.g., bisoprolol, carvedilol, metoprolol) OR
 - 2. Patient has an intolerance, FDA labeled contraindications, or hypersensitivity to a beta blocker

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Cresemba PA

Drug Name(s)

Cresemba

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of invasive aspergillosis OR
 - B. Patient has a diagnosis of invasive mucormycosis OR
 - C. Patient has another indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of invasive aspergillosis and patient has continued indicators of active disease (e.g., continued radiologic findings, direct microscopy findings, histopathology findings, positive cultures, positive serum galactomannan assay) OR
 - B. Patient has a diagnosis of invasive mucormycosis and patient has continued indicators of active disease (e.g., continued radiologic findings, direct microscopy findings, histopathology findings, positive cultures, positive serum galactomannan assay) OR
 - C. BOTH of the following:
 - i. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 6 months

Crysvita PA

Drug Name(s)

Crysvita

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. Patient has a diagnosis of X-linked hypophosphatemia (XLH) as confirmed by ONE of the following:
 - a. Genetic testing OR
 - b. Elevated levels of intact fibroblast growth factor 23 (FGF23) OR
 - c. Prescriber has provided information indicating the patient has a positive family history of XLH AND
 - ii. ONE of the following:
 - a. Patient's epiphyseal plate has not fused OR
 - b. Patient's epiphyseal plate has fused AND the patient is experiencing symptoms of XLH (e.g., bone pain, fractures, limited mobility) OR
 - B. Patient has a diagnosis of tumor-induced osteomalacia (TIO) associated with phosphaturic mesenchymal tumors AND BOTH of the following:
 - i. The requested agent is being used to treat FGF23 related hypophosphatemia AND
 - ii. The tumor cannot be curatively surgically resected or localized AND
- 2. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of X-linked hypophosphatemia (XLH) OR
 - B. Patient has a diagnosis of tumor-induced osteomalacia (TIO) associated with phosphaturic mesenchymal tumors AND
- 3. Patient has had clinical benefit with the requested agent (e.g., enhanced height velocity, improvement in lower extremity bowing and associated abnormalities, radiographic evidence of epiphyseal healing, improvement in bone pain, enhanced mobility, improvement in osteomalacia, improvement in fracture healing) AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Patient is within the FDA labeled age for the requested agent for the requested indication

Prescriber is a specialist in the area of the patient's diagnosis (e.g., nephrologist, endocrinologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Cystadrops PA

Drug Name(s)

Cystadrops

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., ophthalmologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Cystaran PA

Drug Name(s)

Cystaran

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., ophthalmologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Cystinosis Agents PA – Cystagon

Drug Name(s)

Cystagon

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of nephropathic cystinosis AND
- 2. Prescriber has performed a baseline white blood cell (WBC) cystine level test AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of nephropathic cystinosis AND
- 3. Patient has had clinical benefit with the requested agent (e.g., decrease in WBC cystine levels from baseline) AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Dalfampridine PA

Drug Name(s)

Dalfampridine Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has a diagnosis of multiple sclerosis (MS) AND
- 2. ONE of the following:
 - A. The requested agent will be used in combination with a disease modifying agent [e.g., Aubagio, Avonex, Bafiertam, Betaseron, dimethyl fumarate (e.g., Tecfidera), Extavia, fingolimod, glatiramer (e.g., Copaxone, Glatopa), Kesimpta, Mavenclad, Mayzent, Plegridy, Ponvory, Rebif, Vumerity, Zeposia] OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to a disease modifying agent OR
 - C. Prescriber has provided information indicating that a disease modifying agent is not clinically appropriate for the patient

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of multiple sclerosis (MS) AND
- 3. ONE of the following:
 - A. The requested agent will be used in combination with a disease modifying agent [e.g., Aubagio, Avonex, Bafiertam, Betaseron, dimethyl fumarate (e.g., Tecfidera), Extavia, fingolimod, glatiramer (e.g., Copaxone, Glatopa), Kesimpta, Mavenclad, Mayzent, Plegridy, Ponvory, Rebif, Tecfidera, Vumerity, Zeposia] OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to a disease modifying agent OR
 - C. Prescriber has provided information indicating that a disease modifying agent is not clinically appropriate for the patient AND
- 4. Patient has had improvements or stabilization from baseline in timed walking speed (timed 25-foot walk)

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Initial approval will be for 3 months, renewal approval will be for 12 months

Dayvigo PA

Drug Name(s)

Dayvigo

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Droxidopa PA

Drug Name(s)

Droxidopa

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of neurogenic orthostatic hypotension (nOH) AND
- 2. Prescriber has performed baseline blood pressure readings while the patient is sitting or supine (lying face up), AND also within three minutes of standing from a supine position AND
- 3. Patient has a decrease of at least 20 mmHg in systolic blood pressure or 10 mmHg diastolic blood pressure within three minutes after standing AND
- 4. Patient has persistent and consistent symptoms of neurogenic orthostatic hypotension (nOH) caused by ONE of the following:
 - A. Primary autonomic failure [Parkinson's disease (PD), multiple system atrophy, or pure autonomic failure] OR
 - B. Dopamine beta-hydroxylase deficiency OR
 - C. Non-diabetic autonomic neuropathy AND
- 5. Prescriber has assessed the severity of the patient's baseline symptoms of dizziness, lightheadedness, feeling faint, or feeling like the patient may black out AND
- 6. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of neurogenic orthostatic hypotension (nOH) AND
- 3. Patient has had improvements or stabilization with the requested agent as indicated by improvement in severity from baseline symptoms of ONE of the following:
 - A. Dizziness
 - B. Lightheadedness
 - C. Feeling faint
 - D. Feeling like the patient may black out AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist, neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be 1 month for initial, 3 months for renewal

Dupixent PA

Drug Name(s)

Dupixent

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of moderate-to-severe atopic dermatitis AND ALL of the following:
 - i. ONE of the following:
 - a. Patient has tried and failed a topical steroid (e.g., triamcinolone) OR
 - b. Patient has an intolerance, hypersensitivity, or an FDA labeled contraindication to a topical steroid AND
 - ii. For patients 2 years of age or over, ONE of the following:
 - a. Patient has tried and failed a topical calcineurin inhibitor (e.g., pimecrolimus, tacrolimus) OR
 - b. Patient has an intolerance, hypersensitivity, or an FDA labeled contraindication to a topical calcineurin inhibitor AND
 - iii. Patient will NOT be using the requested agent in combination with another biologic agent or a JAK inhibitor for the requested indication (e.g., Adbry, Cibinqo, Opzelura, Rinvoq) OR
 - B. Patient has a diagnosis of moderate-to-severe asthma with an eosinophilic phenotype or oral corticosteroid dependent asthma AND BOTH of the following:
 - i. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LRTA, LAMA, theophylline) in combination with the requested agent AND ii. Patient will NOT be using the requested agent in combination with Xolair or with an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication OR
 - C. Patient has a diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) AND the following:
 - i. BOTH of the following:
 - a. ONE of the following:
 - 1. Patient has tried and had an inadequate response to an oral systemic corticosteroid AND an intranasal corticosteroid (e.g., fluticasone) OR
 - 2. Patient has an intolerance, hypersensitivity, or an FDA labeled contraindication to an oral systemic corticosteroid AND an intranasal corticosteroid AND

Initial criteria continues: see Other Criteria

Age Restriction:

For diagnosis of moderate-to-severe atopic dermatitis, patient is 6 months of age or over. For diagnosis of moderate-to-severe asthma with an eosinophilic phenotype or oral corticosteroid dependent asthma, patient is 6 years of age or over. For diagnosis of CRSwNP, patient is 18 years of age or over. For diagnosis of EoE, patient is 1 year of age or over. For diagnosis of PN, patient is 18 years of age or over.

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, dermatologist, immunologist, gastroenterologist, otolaryngologist, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- b. Patient will continue standard maintenance therapy (e.g., intranasal corticosteroid) in combination with the requested agent OR
- D. Patient has a diagnosis of eosinophilic esophagitis (EoE) confirmed by esophageal biopsy OR
- E. Patient has a diagnosis of prurigo nodularis (PN) AND
- 2. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of moderate-to-severe atopic dermatitis AND the following:
 - i. Patient will NOT be using the requested agent in combination with another biologic agent or a JAK inhibitor for the requested indication (e.g., Adbry, Cibinqo, Opzelura, Rinvoq) OR
 - B. Patient has a diagnosis of moderate-to-severe asthma with an eosinophilic phenotype or oral corticosteroid dependent asthma AND BOTH of the following:
 - i. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LTRA, LAMA, theophylline) in combination with the requested agent AND ii. Patient will NOT be using the requested agent in combination with Xolair or with an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication OR
 - C. Patient has a diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) AND the following:
 - i. Patient will continue standard maintenance therapy (e.g., intranasal corticosteroid) in combination with the requested agent OR
 - D. Patient has a diagnosis of eosinophilic esophagitis (EoE) OR
 - E. Patient has a diagnosis of prurigo nodularis (PN) AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Emgality PA

Drug Name(s)

Emgality

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of migraine AND ALL of the following:
 - i. The requested agent is being used for migraine prophylaxis AND
 - ii. Patient has 4 migraine headaches or more per month AND
 - iii. ONE of the following:
 - a. Patient has tried and had an inadequate response to a conventional migraine prophylaxis agent [e.g., beta blockers (propranolol), anticonvulsants (divalproex, topiramate)] OR
 - b. Patient has an intolerance, or hypersensitivity to a conventional migraine prophylaxis agent OR
 - c. Patient has an FDA labeled contraindication to a conventional migraine prophylaxis agent AND
 - iv. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis OR
 - B. Patient has a diagnosis of episodic cluster headache AND BOTH of the following:
 - i. Patient has had at least 5 cluster headache attacks AND
 - ii. Patient has had at least two cluster periods lasting 7 days to one year and separated by pain-free remission periods of 3 months or more

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. ALL of the following:
 - i. Patient has a diagnosis of migraine AND
 - ii. The requested agent is being used for migraine prophylaxis AND
 - iii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis OR
 - B. Patient has a diagnosis of episodic cluster headache AND
- 3. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Emsam PA

Drug Name(s)

Emsam

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of major depressive disorder (MDD) OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - a. BOTH of the following:
 - i. Patient has a diagnosis of major depressive disorder (MDD) AND
 - ii. ONE of the following:
 - 1. Patient has tried and had an inadequate response to at least two different oral antidepressants (e.g., SSRIs, SNRIs, mirtazapine, bupropion) OR
 - 2. Patient has an intolerance or hypersensitivity to at least two different oral antidepressants (e.g., SSRIs, SNRIs, mirtazapine, bupropion) OR
 - 3. Patient has an FDA labeled contraindication to at least two different oral antidepressants (e.g., SSRIs, SNRIs, mirtazapine, bupropion) OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:

- A. Patient has a diagnosis of major depressive disorder (MDD) OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days

OF

- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. BOTH of the following:
 - i. Patient has had clinical benefit with the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Endari PA

Drug Name(s)

Endari

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of sickle cell disease AND
- 2. Patient is using the requested agent to reduce the acute complications of sickle cell disease AND
- 3. ONE of the following:
 - A. Patient has tried and had an inadequate response to maximally tolerated dose of hydroxyurea OR
 - B. Patient has an intolerance or hypersensitivity to hydroxyurea OR
 - C. Patient has an FDA labeled contraindication to hydroxyurea AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Epclusa PA

Drug Name(s)

Epclusa

Sofosbuvir/Velpatasvir

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of hepatitis C confirmed by serological markers OR
 - B. Patient is a hepatitis C virus (HCV) uninfected solid organ transplant recipient AND BOTH of the following:
 - i. Patient received an HCV viremic donor organ AND
 - ii. The requested agent is being used for prophylaxis AND
- 2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
- 3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
- 4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND
- 5. ONE of the following:
 - A. The requested agent is the preferred agent: Epclusa OR
 - B. The requested agent is the non-preferred agent: sofosbuvir/velpatasvir AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days $\sf OR$
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR
 - iv. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Epidiolex PA

Drug Name(s)

Epidiolex

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of seizures associated with ONE of the following:
 - A. Lennox-Gastaut syndrome OR
 - B. Dravet syndrome OR
 - C. Tuberous sclerosis complex AND
- 2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Erythropoietin Stimulating Agents PA – Aranesp

Drug Name(s)

Aranesp Albumin Free

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. The requested agent is being prescribed for ONE of the following:
 - A. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of the following:
 - i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND
 - ii. Patient is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND
 - iii. The intent of chemotherapy is non-curative OR
 - B. Anemia associated with chronic kidney disease in a patient NOT on dialysis AND ALL of the following:
 - i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks) AND
 - ii. The rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND
 - iii. The intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR
 - C. Anemia due to myelodysplastic syndrome AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) OR
 - D. Another indication that is supported in CMS approved compendia for the requested agent AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND
- 2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

6 months for chemotherapy, 12 months for other indications

Erythropoietin Stimulating Agents PA - Epogen/Procrit

Drug Name(s)

Procrit

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. The requested agent is being prescribed for ONE of the following:
 - A. To reduce the possibility of allogeneic blood transfusion in a surgery patient AND the patient's hemoglobin level is greater than 10 g/dL but 13 g/dL or less OR
 - B. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of the following:
 - i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND
 - ii. Patient is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND
 - iii. The intent of chemotherapy is non-curative OR
 - C. Anemia associated with chronic kidney disease in a patient NOT on dialysis AND ALL of the following:
 - i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks) AND
 - ii. The rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND
 - iii. The intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR
 - D. Anemia due to myelodysplastic syndrome AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) OR
 - E. Anemia resulting from zidovudine treatment of HIV infection AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

1 month for surgery (reduce transfusion possibility), 6 months for chemo, 12 months for other

Other Criteria:

F. Another indication that is supported in CMS approved compendia for the requested agent AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Erythropoietin Stimulating Agents PA – Retacrit

Drug Name(s)

Retacrit

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. The requested agent is being prescribed for ONE of the following:
 - A. To reduce the possibility of allogeneic blood transfusion in a surgery patient AND the patient's hemoglobin level is greater than 10 g/dL but 13 g/dL or less OR
 - B. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of the following:
 - i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND
 - ii. Patient is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND
 - iii. The intent of chemotherapy is non-curative OR
 - C. Anemia associated with chronic kidney disease in a patient NOT on dialysis AND ALL of the following:
 - i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks)

AND

- ii. The rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion
- iii. The intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR
- D. Anemia resulting from zidovudine treatment of HIV infection AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) OR
- E. Another indication that is supported in CMS approved compendia for the requested agent AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND
- 2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

1 month for surgery (reduce transfusion possibility), 6 months for chemo, 12 months for other **Other Criteria:**

Eysuvis PA

Drug Name(s)

Eysuvis

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of dry eye disease AND
- 2. The requested agent will be used for short-term (up to two weeks) treatment AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 1 month

Fasenra PA

Drug Name(s)

Fasenra

Fasenra Pen

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of severe asthma with an eosinophilic phenotype AND
- 2. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LTRA, LAMA, theophylline) in combination with the requested agent AND
- 3. Patient will NOT be using the requested agent in combination with Xolair, Dupixent, or with another injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Nucala) for the requested indication AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of severe asthma with an eosinophilic phenotype AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LTRA, LAMA, theophylline) in combination with the requested agent AND
- 5. Patient will NOT be using the requested agent in combination with Xolair, Dupixent, or with another injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Nucala) for the requested indication AND
- 6. The requested dose is within the FDA labeled dosing for the requested indication

Age Restriction:

Patient is 12 years of age or over

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, immunologist, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Fentanyl Oral PA - Fentanyl lozenge

Drug Name(s)

Fentanyl Citrate Oral Transmucosal

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. ONE of the following:
 - a. Patient has a documented diagnosis (i.e., medical records) of chronic cancer pain due to an active malignancy AND the following:
 - i. There is evidence of a claim that the patient is currently being treated with a longacting opioid with the requested agent within the past 90 days OR
 - b. Patient has a diagnosis that is supported in CMS approved compendia for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with any other oral or nasal fentanyl agent

Age Restriction:

Patient is 16 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Fintepla PA

Drug Name(s)

Fintepla

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of seizures associated with Dravet syndrome (DS) or Lennox-Gastaut syndrome (LGS) AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. An echocardiogram assessment will be obtained before and during treatment with the requested agent, to evaluate for valvular heart disease and pulmonary arterial hypertension AND
 - ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Focalin PA

Drug Name(s)

Dexmethylphenidate Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Gammagard/Gammaked/Gamunex-C PA

Drug Name(s)

Gammagard Liquid

Gammagard S/D Iga Less Than 1Mcg/Ml

Gamunex-C

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ONE of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Primary immunodeficiency [e.g., congenital agammaglobulinemia, common variable immunodeficiency (CVID), severe combined immunodeficiency, Wiskott-Aldrich Syndrome, X-linked agammaglobulinemia (XLA), humoral immunodeficiency, IgG subclass deficiency with or without IgA deficiency] OR
 - B. B-cell chronic lymphocytic leukemia OR multiple myeloma AND ONE of the following:
 - i. Patient has a history of infections OR
 - ii. Patient has evidence of specific antibody deficiency OR
 - iii. Patient has hypogammaglobulinemia OR
 - C. Idiopathic thrombocytopenia purpura AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone), or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - D. Dermatomyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - E. Polymyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - F. Severe rheumatoid arthritis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., tumor necrosis factor antagonists (e.g., Humira), DMARDS (e.g., methotrexate)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 6 months for indications in Other Criteria, 12 months for all others **Other Criteria**:

- G. Myasthenia gravis (MG) AND ONE of the following:
 - i. Patient is in acute myasthenic crisis OR
 - ii. Patient has severe refractory MG (e.g., major functional disability/weakness) AND ONE of the following:
 - a) Patient has failed ONE immunomodulator therapy (i.e., corticosteroid, pyridostigmine, or azathioprine) OR
 - b) Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE immunomodulator therapy OR
- H. Multiple sclerosis (MS) AND BOTH of the following:
 - i. Patient has a diagnosis of relapsing remitting MS (RRMS) AND
 - ii. Patient has had an insufficient response, documented failure, or FDA labeled contraindication to TWO MS agents (e.g., Avonex, Betaseron, Copaxone, dimethyl fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR
- I. Acquired von Willebrand hemophilia AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g., desmopressin solution, von Willebrand factor replacement therapy, corticosteroids, cyclophosphamide, FEIBA, or recombinant factor VIIa) OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- J. Refractory pemphigus vulgaris AND ONE of the following:
 - i. Patient has failed ONE conventional immunosuppressive therapy (e.g., azathioprine, cyclophosphamide, mycophenolate, corticosteroids) OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional immunosuppressive therapy OR
- 2. ONE of the following:
 - A. Patient has another FDA labeled indication for the requested agent OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Indications with 6 months approval duration: Acquired von Willebrand hemophilia, Guillain-Barre Syndrome, Lambert-Eaton myasthenia syndrome, Kawasaki disease, CMV induced pneumonitis in solid organ transplant, Toxic shock syndrome due to invasive group A streptococcus, Toxic epidermal necrolysis and Stevens-Johnson syndrome

Drug is also subject to Part B versus Part D review.

Gattex PA

Drug Name(s)

Gattex

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of short bowel syndrome (SBS) AND
- 2. Patient is dependent on parenteral nutrition OR intravenous (PN/IV) fluids AND
- 3. ONE of the following:
 - A. Patient is aged 1 year to 17 years AND BOTH of the following:
 - i. A fecal occult blood test has been performed within 6 months prior to initiating treatment with the requested agent AND
 - ii. ONE of the following:
 - a. There was no unexplained blood in the stool OR
 - b. There was unexplained blood in the stool AND a colonoscopy or a sigmoidoscopy was performed OR
 - B. Patient is 18 years of age or over AND BOTH of the following:
 - i. Patient has had a colonoscopy within 6 months prior to initiating treatment with the requested agent AND
 - ii. If polyps were present at this colonoscopy, the polyps were removed AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of short bowel syndrome (SBS) AND
- 3. Patient has had a reduction from baseline in parenteral nutrition OR intravenous (PN/IV) fluids AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal

Gaucher Enzyme Replacement PA - Cerezyme

Drug Name(s)

Cerezyme

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of Gaucher disease type 1 (GD1) confirmed by ONE of the following:
 - A. A baseline glucocerebrosidase enzyme activity of less than or equal to 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR
 - B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND
- 2. Prescriber has drawn baseline measurements of hemoglobin, platelet count, liver volume, and spleen volume AND
- 3. Prior to any treatment for the intended diagnosis, the patient has had at least ONE of the following clinical presentations:
 - A. Anemia [defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender] OR
 - B. Thrombocytopenia (defined as a platelet count of less than 100,000 per microliter) OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth failure (i.e., growth velocity below the standard mean for age) OR
 - F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of Gaucher disease type 1 (GD1) AND
- 3. Patient has had improvement and/or stabilization from baseline in at least ONE of the following:
 - A. Hemoglobin (Hb) level OR
 - B. Platelet count OR
 - C. Liver volume OR
 - D. Spleen volume OR
 - E. Growth velocity OR
 - F. Bone pain or disease

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Gaucher Enzyme Replacement PA – Elelyso

Drug Name(s)

Elelyso

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of Gaucher disease type 1 (GD1) confirmed by ONE of the following:
 - A. A baseline glucocerebrosidase enzyme activity of less than or equal to 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR
 - B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND
- 2. Prescriber has drawn baseline measurements of hemoglobin, platelet count, liver volume, and spleen volume AND
- 3. Prior to any treatment for the intended diagnosis, the patient has had at least ONE of the following clinical presentations:
 - A. Anemia [defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender] OR
 - B. Thrombocytopenia (defined as a platelet count of less than 100,000 per microliter) OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth failure (i.e., growth velocity below the standard mean for age) OR
 - F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of Gaucher disease type 1 (GD1) AND
- 3. Patient has had improvement and/or stabilization from baseline in at least ONE of the following:
 - A. Hemoglobin (Hb) level OR
 - B. Platelet count OR
 - C. Liver volume OR
 - D. Spleen volume OR
 - E. Growth velocity OR
 - F. Bone pain or disease

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Gaucher Enzyme Replacement PA - Vpriv

Drug Name(s)

Vpriv

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of Gaucher disease type 1 (GD1) confirmed by ONE of the following:
 - A. A baseline glucocerebrosidase enzyme activity of less than or equal to 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR
 - B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND
- 2. Prescriber has drawn baseline measurements of hemoglobin, platelet count, liver volume, and spleen volume AND
- 3. Prior to any treatment for the intended diagnosis, the patient has had at least ONE of the following clinical presentations:
 - A. Anemia [defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender] OR
 - B. Thrombocytopenia (defined as a platelet count of less than 100,000 per microliter) OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth failure (i.e., growth velocity below the standard mean for age) OR
 - F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of Gaucher disease type 1 (GD1) AND
- 3. Patient has had improvement and/or stabilization from baseline in at least ONE of the following:
 - A. Hemoglobin (Hb) level OR
 - B. Platelet count OR
 - C. Liver volume OR
 - D. Spleen volume OR
 - E. Growth velocity OR
 - F. Bone pain or disease

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Growth Hormone PA – Omnitrope

Drug Name(s)

Omnitrope

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

For Children – Criteria for initial approval require the following:

- 1. ONE of the following:
 - a. Patient has a diagnosis of Turner Syndrome OR
 - b. Patient has a diagnosis of Prader-Willi Syndrome OR
 - c. Patient has a diagnosis of panhypopituitarism AND BOTH of the following:
 - i. Deficiencies in 3 or more pituitary axes AND
 - ii. Measured serum IGF-1 (insulin-like growth factor-1) levels are below the age and sexappropriate reference range when off GH therapy OR
 - d. Patient has a diagnosis of growth hormone deficiency (GHD) or short stature AND BOTH of the following:
 - i. Patient has ONE of the following:
 - a) Height more than 2 standard deviations (SD) below the mean for age and sex OR
 - b) Height more than 1.5 SD below the midparental height OR
 - c) A decrease in height SD of more than 0.5 over one year in children at least 2 years of age OR
 - d) Height velocity more than 2 SD below the mean over one year or more than 1.5 SD sustained over two years AND
 - ii. Failure of at least 2 growth hormone (GH) stimulation tests (e.g., peak GH value of less than 10 mcg/L after stimulation, or otherwise considered abnormal as determined by testing lab) OR
 - e. Patient has a diagnosis of small for gestational age (SGA) AND ALL of the following:
 - i. Patient is at least 2 years of age AND
 - ii. Documented birth weight and/or length that is 2 or more SD below the mean for gestational age AND
 - iii. At 24 months of age, the patient fails to manifest catch-up growth evidenced by a height that remains 2 or more SD below the mean for age and sex

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

For Children – Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the preferred agent through the plan's Prior Authorization criteria AND
- 2. Patient has been diagnosed with ONE of the following:
 - a. Growth Hormone Deficiency, Short Stature OR
 - b. Panhypopituitarism OR
 - c. Prader-Willi Syndrome OR
 - d. Small for Gestational Age (SGA) OR
 - e. Turner Syndrome AND
- 3. ALL of the following:
 - a. Patient does NOT have closed epiphyses AND
 - b. Patient is being monitored for adverse effects of therapy with the requested agent AND
 - c. Patient's height has increased or height velocity has improved since initiation or last approval of the requested agent

For Adults – Criteria for initial approval require the following:

- 1. Patient has been diagnosed with ONE of the following:
 - a. Childhood growth hormone deficiency (GHD) with genetic or organic origin AND ONE of the following:
 - i. Low IGF-1 (insulin-like growth factor-1) level without GH replacement therapy OR
 - ii. Failure of at least one growth hormone (GH) stimulation test as an adult (e.g., peak GH value of 5 mcg/L or lower after stimulation, or otherwise considered abnormal as determined by testing lab) OR
 - b. Acquired adult GHD secondary to structural lesions or trauma AND ONE of the following:
 - i. Patient has a diagnosis of panhypopituitarism AND BOTH of the following:
 - a) Deficiencies in 3 or more pituitary axes AND
 - b) Low IGF-1 level without GH replacement therapy OR
 - ii. Patient has failed at least one growth hormone (GH) stimulation test as an adult OR c. Idiopathic GHD (adult or childhood onset) AND the patient has failed at least two growth hormone (GH) stimulation tests as an adult

For Adults – Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the preferred agent through the plan's Prior Authorization criteria AND
- 2. Patient has been diagnosed with ONE of the following:
 - a. Childhood growth hormone deficiency (GHD) with genetic or organic origin OR
 - b. Acquired adult GHD secondary to structural lesions or trauma OR
 - c. Idiopathic GHD (adult or childhood onset) AND
- 3. Patient is being monitored for adverse effects of therapy with the requested agent AND
- 4. Patient's IGF-1 level has been evaluated to confirm the appropriateness of the current dose AND
- 5. Patient has had clinical benefit with the requested agent (i.e., body composition, hip-to-waist ratio, cardiovascular health, bone mineral density, serum cholesterol, physical strength, or quality of life)

HAE PA - Cinryze

Drug Name(s)

Cinryze

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Acute HAE attacks

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of hereditary angioedema (HAE) which has been confirmed via two confirmatory tests of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:
 - a. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type I)]: decreased quantities of C4 and C1-INH (antigenic and functional level) OR
 - b. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type II)]: decreased quantities of C4 and C1-INH function (C1-INH protein level may be normal) OR
 - c. Hereditary angioedema (HAE) with normal C1INH [HAE-nI-C1INH (Type III)]: Normal levels of C4 and C1-INH [antigenic and functional level (at baseline and during an attack)] AND ONE of the following:
 - i. BOTH of the following:
 - 1. Family history of angioedema AND
 - 2. ALL other causes of angioedema have been ruled out OR
 - ii. Patient demonstrates a Factor XII mutation, angiopoietin-1 (ANGPT1) mutation, plasminogen (PLG) mutation, kininogen1 mutation, heparan sulfate 3-O-sulfotransferase 6 gene mutation, or myoferlin gene mutation that is associated with the disease AND
- 2. Medications known to cause angioedema (e.g., ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND
- 3. ONE of the following:
 - a. The requested agent will be used to treat acute HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks OR
 - b. The requested agent will be used for prophylaxis against HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria

AND

- 2. Patient has a diagnosis of hereditary angioedema (HAE) AND ONE of the following:
 - a. The requested agent will be used to treat acute HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks OR
 - b. The requested agent will be used for prophylaxis against HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks AND
- 3. Patient has had a decrease in the frequency or severity of acute attacks or has had stabilization of disease from use of the requested agent

HAE PA - Haegarda

Drug Name(s)

Haegarda

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of hereditary angioedema (HAE) which has been confirmed via two confirmatory tests of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:
 - a. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type I)]: decreased quantities of C4 and C1-INH (antigenic and functional level) OR
 - b. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type II)]: decreased quantities of C4 and C1-INH function (C1-INH protein level may be normal) OR
 - c. Hereditary angioedema (HAE) with normal C1INH [HAE-nI-C1INH (Type III)]: Normal levels of C4 and C1-INH [antigenic and functional level (at baseline and during an attack)] AND ONE of the following:
 - i. BOTH of the following:
 - 1. Family history of angioedema AND
 - 2. ALL other causes of angioedema have been ruled out OR
 - ii. Patient demonstrates a Factor XII mutation, angiopoietin-1 (ANGPT1) mutation, plasminogen (PLG) mutation, kininogen1 mutation, heparan sulfate 3-O-sulfotransferase 6 gene mutation, or myoferlin gene mutation that is associated with the disease AND
- 2. Medications known to cause angioedema (e.g., ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND
- 3. The requested agent will be used for prophylaxis against HAE attacks AND
- 4. Patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of hereditary angioedema (HAE) AND
- 3. The requested agent is being used for prophylaxis against HAE attacks AND
- 4. Patient has had a decrease in the frequency or severity of acute attacks or has had stabilization of disease from use of the requested agent AND

5. Patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks

HAE PA – Icatibant

Drug Name(s)

Icatibant Acetate

Sajazir

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of hereditary angioedema (HAE) which has been confirmed via two confirmatory tests of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:
 - a. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type I)]: decreased quantities of C4 and C1-INH (antigenic and functional level) OR
 - b. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type II)]: decreased quantities of C4 and C1-INH function (C1-INH protein level may be normal) OR
 - c. Hereditary angioedema (HAE) with normal C1INH [HAE-nI-C1INH (Type III)]: Normal levels of C4 and C1-INH [antigenic and functional level (at baseline and during an attack)] AND ONE of the following:
 - i. BOTH of the following:
 - 1. Family history of angioedema AND
 - 2. ALL other causes of angioedema have been ruled out OR
 - ii. Patient demonstrates a Factor XII mutation, angiopoietin-1 (ANGPT1) mutation, plasminogen (PLG) mutation, kininogen1 mutation, heparan sulfate 3-O-sulfotransferase 6 gene mutation, or myoferlin gene mutation that is associated with the disease AND
- 2. Medications known to cause angioedema (e.g., ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND
- 3. The requested agent will be used to treat acute HAE attacks AND
- 4. Patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of hereditary angioedema (HAE) AND
- 3. The requested agent will be used to treat acute HAE attacks AND

- 4. Patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks AND
- 5. Patient has had a decrease in the frequency or severity of acute attacks or stabilization of disease from use of the requested agent

Harvoni PA

Drug Name(s)

Harvoni

Ledipasvir/Sofosbuvir

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND
- 2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
- 3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
- 4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA dosing for the requested indication AND
- 5. ONE of the following:
 - A. The requested agent is the preferred agent: Harvoni OR
 - B. The requested agent is the non-preferred agent: ledipasvir/sofosbuvir AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past $90\ \text{days}\ \text{OR}$
 - iii. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR
 - iv. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported **Other Criteria**:

High Risk Medication PA - All Starts

Drug Name(s)

Benztropine Mesylate

Clemastine Fumarate

Cyproheptadine Hydrochloride

Dicyclomine Hydrochloride

Diphenoxylate Hydrochloride/Atropine Sulfate

Hydroxyzine Hydrochloride

Hydroxyzine Pamoate

Methscopolamine Bromide

Promethazine Hcl Plain

Promethazine Hydrochloride

Promethegan

Scopolamine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

PA does NOT apply to patients less than 65 years of age.

Criteria for approval require ALL of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested high-risk medication AND
- 2. Prescriber has indicated that the benefits of the requested high-risk medication outweigh the risks for the patient AND
- 3. Prescriber has indicated that the risks and potential side effects of the requested high-risk medication have been discussed with the patient

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Ilaris PA

Drug Name(s)

Ilaris

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Acute gouty arthritis

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of:

- 1. ONE of the following:
 - A. Patient (pt) has been diagnosed with Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS) OR
 - B. Pt has been diagnosed with Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD) OR
 - C. Pt has been diagnosed with Familial Mediterranean Fever (FMF) AND ONE of the following:
 - i. Pt has tried and had an inadequate response to colchicine OR
 - ii. Pt has an intolerance or hypersensitivity to colchicine OR
 - iii. Pt has an FDA labeled contraindication to colchicine OR
 - D. Pt has been diagnosed with Cryopyrin-Associated Periodic Syndrome (CAPS) including Familial Cold Auto-inflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS) OR
 - E. Pt has been diagnosed with active Systemic Juvenile Idiopathic Arthritis (SJIA) AND BOTH of the following:
 - i. Pt has documented active systemic features (e.g., ongoing fever, evanescent erythematous rash, generalized lymphadenopathy, 1 or more joints with active arthritis, hepatomegaly, splenomegaly, serositis) AND
 - ii. ONE of the following:
 - a. Pt has tried and had an inadequate response to at least ONE prerequisite agent (e.g., glucocorticosteroids, prescription oral NSAIDs, methotrexate, leflunomide) OR
 - b. Pt has an intolerance or hypersensitivity to at least ONE prerequisite agent OR
 - c. Pt has an FDA labeled contraindication to at least ONE prerequisite agent OR
 - F. Pt has a diagnosis of adult onset Still's disease OR
 - G. Pt has been diagnosed with acute gouty arthritis AND ONE of the following:
 - i. Pt has tried and had an inadequate response to at least TWO conventional first-line agents (e.g., prescription oral NSAIDs, colchicine, systemic corticosteroids) OR
 - ii. Pt has an intolerance or hypersensitivity to at least TWO conventional first-line agents OR
 - iii. Pt has an FDA labeled contraindication to at least TWO conventional first-line agents AND
- 2. Pt will NOT be using the requested agent in combination with another biologic agent **Age Restriction:**

For diagnosis of CAPS including FCAS or MWS, patient is 4 years of age or over

For diagnosis of SJIA, patient is 2 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Imiquimod PA

Drug Name(s)

Imiquimod

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Actinic keratosis OR
 - B. Superficial basal cell carcinoma OR
 - C. External genital and/or perianal warts/condyloma acuminata OR
 - D. Squamous cell carcinoma OR
 - E. Basal cell carcinoma OR
 - F. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

4 months for Actinic keratosis, other diagnoses - see Other Criteria

Other Criteria:

2 months for Superficial basal cell carcinoma, Squamous cell carcinoma, or Basal cell carcinoma

4 months for External genital and/or perianal warts/condyloma acuminata

12 months for All other diagnoses

Inbrija PA

Drug Name(s)

Inbrija

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. The requested agent will be used for intermittent treatment of OFF episodes in patients with Parkinson's disease AND
- 2. Patient is receiving concurrent therapy with carbidopa/levodopa AND
- 3. Patient will NOT be using a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine, tranylcypromine) in combination with, or within 2 weeks of, the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Injectable Oncology PA Drug Name(s) Abraxane Adcetris Aliqopa Alymsys Avastin Beleodaq Besponsa Blenrep Blincyto Bortezomib Cyramza Danyelza Darzalex Darzalex Faspro Doxorubicin Hydrochloride Liposomal Empliciti Enhertu Erbitux Folotyn Fulvestrant Gazyva Halaven Herceptin Herceptin Hylecta Herzuma Jevtana Kadcyla Kanjinti **Kyprolis** Lumoxiti Margenza Monjuvi Mvasi

Mylotarg

Nelarabine
Ogivri
Onivyde
Ontruzant
Padcev
Pemetrexed
Perjeta
Phesgo
Polivy
Portrazza
Poteligeo
Romidepsin
Rybrevant
Sarclisa
Synribo
Trazimera
Trodelvy
Unituxin
Vectibix
Vegzelma
Vyxeos
Yondelis
Zaltrap
Zepzelca
Zirabev
Zynlonta
Indications:
All Medically-Accepted Indications. Off-Label Uses:
Exclusion Criteria:
Required Medical Information:
Criteria for approval require BOTH of the following:
1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia
for the requested agent AND 2. ONE of the following:
A. There is evidence of a claim that the patient is currently being treated with the requested
agent within the past 180 days OR
B. Prescriber states the patient is currently being treated with the requested agent OR
C. ALL of the following:

- i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
- ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported by CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested indication OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
- iii. Patient does NOT have any FDA labeled contraindications to the requested agent AND
- iv. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

May also be subject to Part B versus Part D review.

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Iron Chelating Agents PA - Exjade

Drug Name(s)

Deferasirox (Exjade)

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome AND ONE of the following:
 - i. A liver iron (Fe) concentration (LIC) of at least 5 mg Fe per gram of dry weight OR
 - ii. A serum ferritin greater than 300 mcg/L OR
 - iii. MRI confirmation of iron deposition OR
 - B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND
- 2. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome OR
 - B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent for the requested indication

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Iron Chelating Agents PA – Jadenu

Drug Name(s)

Deferasirox (Jadenu)

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome AND ONE of the following:
 - i. A liver iron (Fe) concentration (LIC) of at least 5 mg Fe per gram of dry weight OR
 - ii. A serum ferritin greater than 300 mcg/L OR
 - iii. MRI confirmation of iron deposition OR
 - B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND
- 2. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome OR
 - B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent for the requested indication

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Ivermectin Cream PA

Drug Name(s)

Ivermectin Cream

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Ivermectin Tablet PA

Drug Name(s)

Ivermectin Tablet

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 4 months

Kalydeco PA

Drug Name(s)

Kalydeco

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of cystic fibrosis AND
- 2. ONE of the following:
 - A. Patient has ONE of the CFTR gene mutations or a mutation in the CFTR gene that is responsive based on in vitro data, as indicated in the FDA label, confirmed by genetic testing OR
 - B. Patient has another CFTR gene mutation(s) that is responsive to the requested agent, as indicated in the FDA label, confirmed by genetic testing AND
- 3. Patient is NOT homozygous for the F508del mutation AND
- 4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of cystic fibrosis AND
- 3. Patient has had improvement or stabilization with the requested agent [e.g., improvement in FEV1 from baseline, increase in weight/BMI, improvement from baseline Cystic Fibrosis Questionnaire-Revised (CFQ-R) Respiratory Domain score, improvements in respiratory symptoms (cough, sputum production, and difficulty breathing), and/or reduced number of pulmonary exacerbations] AND
- 4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Kerendia PA

Drug Name(s)

Kerendia

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Korlym PA

Drug Name(s)

Korlym

Mifepristone

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of Cushing's syndrome AND
- 2. ONE of the following:
 - A. Patient has type 2 diabetes mellitus OR
 - B. Patient has glucose intolerance as defined by a 2-hour glucose tolerance test plasma glucose value of 140-199 mg/dL AND
- 3. ONE of the following:
 - A. Patient has failed surgical resection OR
 - B. Patient is NOT a candidate for surgical resection

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of Cushing's syndrome AND
- 3. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Leuprolide PA

Drug Name(s)

Eligard

Leuprolide Acetate

Lupron Depot (1-Month)

Lupron Depot (3-Month)

Lupron Depot (4-Month)

Lupron Depot (6-Month)

Lupron Depot-Ped (1-Month)

Lupron Depot-Ped (3-Month)

Lupron Depot-Ped (6-Month)

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. BOTH of the following:
 - i. Patient is NOT currently being treated with the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent AND
- 3. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA - Lidocaine Gel/Jelly

Drug Name(s)

Glydo

Lidocaine Hcl Gel

Lidocaine Hcl Jelly

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. The requested agent will be used for ONE of the following:
 - A. Surface anesthesia and lubrication for urethral procedure OR
 - B. Topical treatment for pain of urethritis OR
 - C. Surface anesthesia and lubrication for endotracheal intubation (oral and nasal) OR
 - D. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA - Lidocaine Ointment

Drug Name(s)

Lidocaine Ointment

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. The requested agent will be used for ONE of the following:
 - A. Anesthesia of accessible mucous membranes of the oropharynx OR
 - B. Anesthetic lubricant for intubation OR
 - C. Temporary relief of pain associated with minor burns, including sunburn, abrasions of the skin, and insect bites OR
 - D. Another indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - i. Patient has tried and had an inadequate response to a conventional therapy [e.g., gabapentin, pregabalin, oral prescription NSAID (non-steroidal anti-inflammatory drug)] for the requested indication OR
 - ii. Patient has an intolerance or hypersensitivity to a conventional therapy OR
 - iii. Patient has an FDA labeled contraindication to a conventional therapy

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA - Lidocaine Patch

Drug Name(s)

Lidocan

Lidocan III

Lidocaine Patch

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Pain associated with postherpetic neuralgia (PHN) OR
 - B. Pain associated with diabetic neuropathy OR
 - C. Neuropathic pain associated with cancer, or cancer treatment OR
 - D. Another diagnosis that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. Patient has tried and had an inadequate response to a conventional therapy [e.g., gabapentin, pregabalin, oral prescription NSAID (non-steroidal anti-inflammatory drug)] for the requested indication OR
 - B. Patient has an intolerance or hypersensitivity to a conventional therapy OR
 - C. Patient has an FDA labeled contraindication to a conventional therapy

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA - Lidocaine Solution

Drug Name(s)

Lidocaine Solution

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. The requested agent will be used for ONE of the following:
 - A. Topical anesthesia of accessible mucous membranes of the oral and nasal cavities OR
 - B. Topical anesthesia of accessible mucous membranes of proximal portions of the digestive tract OR
 - C. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA - Lidocaine/prilocaine Cream

Drug Name(s)

Lidocaine/Prilocaine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. The requested agent will be used for ONE of the following:
 - A. Local analgesia on normal intact skin OR
 - B. Topical anesthetic for dermal procedures OR
 - C. Adjunctive anesthesia prior to local anesthetic infiltration in adult male genital skin OR
 - D. Anesthesia for minor procedures on female external genitalia OR
 - E. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA - ZTlido

Drug Name(s)

Ztlido

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Pain associated with postherpetic neuralgia (PHN) OR
 - B. Neuropathic pain associated with cancer, or cancer treatment OR
 - C. Another diagnosis that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. Patient has tried and had an inadequate response to generic lidocaine 5% patch OR
 - B. Patient has an intolerance or hypersensitivity to generic lidocaine 5% patch OR
 - C. Patient has an FDA labeled contraindication to generic lidocaine 5% patch AND
- 3. ONE of the following:
 - A. Patient has tried and had an inadequate response to a conventional therapy [e.g., gabapentin, pregabalin, oral prescription NSAID (non-steroidal anti-inflammatory drug)] for the requested indication OR
 - B. Patient has an intolerance or hypersensitivity to a conventional therapy OR
 - C. Patient has an FDA labeled contraindication to a conventional therapy

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Linezolid PA

Drug Name(s)

Linezolid

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND ONE of the following:
 - a. The requested agent is prescribed by an infectious disease specialist or the prescriber has consulted with an infectious disease specialist on treatment of this patient OR
 - b. Patient has a documented infection due to vancomycin-resistant Enterococcus faecium OR
 - c. Patient has a diagnosis of pneumonia caused by Staphylococcus aureus or Streptococcus pneumoniae AND ONE of the following:
 - i. Patient has a documented infection that is resistant to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole, OR that is resistant to vancomycin OR
 - ii. Patient has an intolerance or hypersensitivity to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
 - iii. Patient has an FDA labeled contraindication to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
 - iv. Patient has an intolerance or hypersensitivity to vancomycin OR
 - v. Patient has an FDA labeled contraindication to vancomycin OR
 - d. Patient has a documented skin and skin structure infection, including diabetic foot infections, caused by Staphylococcus aureus, Streptococcus pyogenes, or Streptococcus agalactiae AND ONE of the following:
 - i. Patient has a documented infection that is resistant to TWO of the following: betalactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole, OR that is resistant to vancomycin at the site of infection OR
 - ii. Patient has an intolerance or hypersensitivity to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
 - iii. Patient has an FDA labeled contraindication to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions: Coverage Duration:

Approval will be for 3 months

Other Criteria:

iv. Patient has an intolerance or hypersensitivity to vancomycin OR

- v. Patient has an FDA labeled contraindication to vancomycin AND
- 2. Patient will NOT be using the requested agent in combination with Sivextro (tedizolid) for the same infection AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Memantine ER PA

Drug Name(s)

Memantine Hydrochloride Er

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

PA does NOT apply to patients greater than or equal to 30 years of age Criteria for approval require the following:

- 1. Patient is younger than 30 years of age AND ONE of the following:
 - A. Patient has a diagnosis of moderate to severe dementia of the Alzheimer's type OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Memantine PA

Drug Name(s)

Memantine Hcl Titration Pak

Memantine Hydrochloride

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

PA does NOT apply to patients greater than or equal to 30 years of age Criteria for approval require the following:

- 1. Patient is younger than 30 years of age AND ONE of the following:
 - A. Patient has a diagnosis of moderate to severe dementia of the Alzheimer's type OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Methylin PA

Drug Name(s)

Methylphenidate Hydrochloride (Methylin)

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Methylphenidate ER Tablet PA

Drug Name(s)

Methylphenidate Hydrochloride Er Tablet

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Migranal PA

Drug Name(s)

Dihydroergotamine Mesylate Spray

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. The requested agent will be used for the treatment of acute migraine with or without aura AND
- 2. ONE of the following:
 - A. Patient has tried and had an inadequate response to TWO triptan agents with differing active ingredients (e.g., sumatriptan, rizatriptan) OR
 - B. Patient has an intolerance or hypersensitivity to TWO triptan agents with differing active ingredients OR
 - C. Patient has an FDA labeled contraindication to TWO triptan agents with differing active ingredients AND
- 3. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, acute CGRP)

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. The requested agent will be used for the treatment of acute migraine with or without aura AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, acute CGRP)

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Modafinil PA

Drug Name(s)

Modafinil

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another target agent (i.e., armodafinil)

Age Restriction:

Patient is 17 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA – Avonex

Drug Name(s)

Avonex

Avonex Pen

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA – Betaseron

Drug Name(s)

Betaseron

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA - Dimethyl Fumarate

Drug Name(s)

Dimethyl Fumarate

Dimethyl Fumarate Starterpack

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA - Fingolimod

Drug Name(s)

Fingolimod

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication AND
- 3. Prescriber has performed an electrocardiogram within 6 months prior to initiating treatment

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA – Glatiramer

Drug Name(s)

Copaxone

Glatiramer Acetate

Glatopa

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA – Mayzent

Drug Name(s)

Mayzent

Mayzent Starter Pack

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA – Plegridy

Drug Name(s)

Plegridy

Plegridy Starter Pack

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA – Tysabri

Drug Name(s)

Tysabri

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. Patient has ONE of the following diagnoses:
 - i. Relapsing form of Multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, or active secondary progressive disease AND ONE of the following:
 - a. Prescriber states the patient has highly active disease OR
 - b. ONE of the following:
 - 1. Patient has tried and had an inadequate response to TWO preferred agents (Avonex, Betaseron, Copaxone, dimethyl fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR
 - 2. Patient has an intolerance or hypersensitivity to TWO preferred agents (Avonex, Betaseron, Copaxone, dimethyl fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR
 - 3. Patient has an FDA labeled contraindication to TWO preferred agents (Avonex, Betaseron, Copaxone, dimethyl fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months for MS, for CD 16 weeks for initial and 12 months renewal **Other Criteria**:

- ii. Moderately to severely active Crohn's disease (CD) AND BOTH of the following:
 - a. ONE of the following:
 - 1. Patient has tried and had an inadequate response to ONE conventional CD therapy (e.g., 6-mercaptopurine, azathioprine, corticosteroids, methotrexate) OR

- 2. Patient has an intolerance or hypersensitivity to ONE conventional CD therapy (e.g., 6-mercaptopurine, azathioprine, corticosteroids, methotrexate) OR
- 3. Patient has an FDA labeled contraindication to ONE conventional CD therapy (e.g., 6-mercaptopurine, azathioprine, corticosteroids, methotrexate) AND
- b. ONE of the following:
 - 1. Patient has tried and had an inadequate response to ONE preferred biologic agent (Humira, Skyrizi, or Stelara) for the treatment of CD OR
 - 2. Patient has an intolerance or hypersensitivity to ONE preferred biologic agent (Humira, Skyrizi, or Stelara) OR
 - 3. Patient has an FDA labeled contraindication to ONE preferred biologic agent (Humira, Skyrizi, or Stelara) AND
- 3. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

MS PA – Vumerity

Drug Name(s)

Vumerity

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Myalept PA

Drug Name(s)

Myalept

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has leptin deficiency associated with a diagnosis of either congenital generalized lipodystrophy (CGL) or acquired generalized lipodystrophy (AGL) AND
- 2. Prescriber has provided the patient's baseline levels for HbA1C, triglycerides, and fasting insulin, measured prior to beginning therapy with the requested agent AND
- 3. Patient also has at least ONE of the complications related to lipodystrophy: diabetes mellitus, hypertriglyceridemia (200 mg/dL or higher), and/or high fasting insulin (30µU/mL or higher) AND
- 4. Patient has tried and had an inadequate response to maximum tolerable dosing of a conventional agent for the additional diagnosis AND
- 5. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has leptin deficiency associated with a diagnosis of either congenital generalized lipodystrophy (CGL) or acquired generalized lipodystrophy (AGL) AND
- 3. Patient has had improvement or stabilization with the requested agent as indicated by change from baseline level of at least ONE of the following:
 - A. HbA1C
 - B. Triglycerides
 - C. Fasting insulin AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Conventional agent examples include:

Hypertriglyceridemia: statins, fenofibrates, Omega-3-Acid Ethyl Esters (generic Lovaza)

Diabetes/high fasting insulin: insulin, sulfonylurea/sulfonylurea combination, metformin/metformin combination

Natpara PA

Drug Name(s)

Natpara

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Increased baseline risk for osteosarcoma (e.g., Paget's disease of bone, unexplained elevations of alkaline phosphatase, hereditary disorders predisposing to osteosarcoma, history of external beam or implant radiation therapy involving the skeleton, pediatric and young adult patients with open epiphyses)

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of hypocalcemia associated with hypoparathyroidism AND
- 2. Patient does NOT have a baseline vitamin D level below the testing laboratory's lower limit of normal AND
- 3. Patient's baseline serum calcium level (albumin-corrected) is above 7.5 mg/dL AND
- 4. Patient will NOT be using the requested agent in combination with alendronate for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of hypocalcemia associated with hypoparathyroidism AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with alendronate for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, nephrologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Nuedexta PA

Drug Name(s)

Nuedexta

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of pseudobulbar affect OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- 2. Patient will NOT be using the requested agent in combination with a monoamine oxidase inhibitor (MAOI) [e.g., Marplan (isocarboxazid), Nardil (phenelzine), and Parnate (tranylcypromine)]

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Nuplazid PA

Drug Name(s)

Nuplazid

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Nurtec PA

Drug Name(s)

Nurtec

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has a diagnosis of migraine AND
- 2. ONE of the following:
 - A. The requested agent is being used for the treatment of acute migraine with or without aura AND BOTH of the following:
 - i. ONE of the following:
 - a. Patient has tried and had an inadequate response to a triptan (e.g., sumatriptan, rizatriptan) agent OR
 - b. Patient has an intolerance, or hypersensitivity to a triptan OR
 - c. Patient has an FDA labeled contraindication to a triptan AND
 - ii. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP) OR
 - B. The requested agent is being used for migraine prophylaxis AND ALL of the following:
 - i. Patient has 4 migraine headaches or more per month AND
 - ii. ONE of the following:
 - a. Patient has tried and had an inadequate response to a conventional migraine prophylaxis agent [e.g., beta blockers (propranolol), anticonvulsants (divalproex, topiramate)] OR
 - b. Patient has an intolerance, or hypersensitivity to a conventional migraine prophylaxis agent OR
 - c. Patient has an FDA labeled contraindication to a conventional migraine prophylaxis agent AND
 - iii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 Months

Other Criteria:

Criteria for renewal require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of migraine AND
- 3. ONE of the following:

- A. The requested agent is being used for the treatment of acute migraine with or without aura AND BOTH of the following:
 - i. Patient has had clinical benefit with the requested agent AND
 - ii. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP) OR
- B. The requested agent is being used for migraine prophylaxis AND BOTH of the following:
 - i. Patient has had clinical benefit with the requested agent AND
 - ii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis

Ocaliva PA

Drug Name(s)

Ocaliva

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of primary biliary cholangitis (PBC) confirmed by at least TWO of the following:
 - A. There is biochemical evidence of cholestasis with an alkaline phosphatase (ALP) elevation
 - B. Presence of antimitochondrial antibody (AMA): a titer greater than 1:80 OR a level that is above the testing laboratory's upper limit of the normal range
 - C. If the AMA is negative or present only in low titer (less than or equal to 1:80), presence of other PBC-specific autoantibodies, including sp100 or gp210
 - D. Histologic evidence of nonsuppurative destruction cholangitis and destruction of interlobular bile ducts AND
- 2. ONE of the following:
 - A. Patient does NOT have cirrhosis OR
 - B. Patient has compensated cirrhosis with NO evidence of portal hypertension AND
- 3. Prescriber has measured the patient's alkaline phosphatase (ALP) level AND total bilirubin level AND
- 4. ONE of the following:
 - A. BOTH of the following:
 - i. Patient has tried and had an inadequate response to ursodiol AND
 - ii. The requested agent will be used in combination with ursodiol OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ursodiol

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of primary biliary cholangitis (PBC) AND
- 3. ONE of the following:
 - A. Patient does NOT have cirrhosis OR
 - B. Patient has compensated cirrhosis with NO evidence of portal hypertension AND
- 4. ONE of the following:
 - A. The requested agent will be used in combination with ursodiol OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ursodiol AND

- 5. Patient has had improvements or stabilization with the requested agent as indicated by BOTH of the following:
 - A. Decrease in alkaline phosphatase (ALP) level from baseline AND
 - B. Total bilirubin is less than or equal to the upper limit of normal (ULN)

Ofev PA

Drug Name(s)

Ofev

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
 - ii. Patient has no known explanation for interstitial lung disease (ILD) or pulmonary fibrosis (e.g., radiation, drugs, metal dusts, sarcoidosis, or any connective tissue disease known to cause ILD) OR
 - B. BOTH of the following:
 - i. Patient has a diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD) AND
 - ii. Patient's diagnosis has been confirmed on high-resolution computed tomography (HRCT) or chest radiography scans OR
 - C. BOTH of the following:
 - i. Patient has a diagnosis of chronic fibrosing interstitial lung disease (ILD) with a progressive phenotype AND
 - ii. Patient's diagnosis has been confirmed on high-resolution computed tomography (HRCT)

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of ONE of the following:
 - A. Idiopathic pulmonary fibrosis (IPF) OR
 - B. Systemic sclerosis-associated interstitial lung disease (SSc-ILD) OR
 - C. Chronic fibrosing interstitial lung disease (ILD) with a progressive phenotype AND
- 3. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., pathologist, pulmonologist, radiologist, rheumatologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA – Bavencio

Drug Name(s)

Bavencio

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested agent OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA – Imfinzi

Drug Name(s)

Imfinzi

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested agent OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA - Jemperli

Drug Name(s)

Jemperli

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested agent OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA - Keytruda

Drug Name(s)

Keytruda

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested agent OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA – Libtayo

Drug Name(s)

Libtayo

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested agent OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA - Opdivo

Drug Name(s)

Opdivo

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested agent OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA - Tecentriq

Drug Name(s)

Tecentriq

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested agent OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA – Yervoy

Drug Name(s)

Yervoy

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested agent OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Opioids ER PA - Buprenorphine Pain

Drug Name(s)

Belbuca

Buprenorphine

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
 - B. Patient has a diagnosis of pain due to sickle cell disease OR
 - C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. ALL of the following:
 - a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - d. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR
 - 3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND
 - e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND
 - f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Opioids ER PA - Fentanyl Patch

Drug Name(s)

Fentanyl

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
 - B. Patient has a diagnosis of pain due to sickle cell disease OR
 - C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. ALL of the following:
 - a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - d. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR
 - 3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND
 - e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND
 - f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Opioids ER PA - Hydrocodone

Drug Name(s)

Hydrocodone Bitartrate Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
 - B. Patient has a diagnosis of pain due to sickle cell disease OR
 - C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. ALL of the following:
 - a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - d. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR
 - 3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND
 - e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND
 - f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Opioids ER PA - Morphine

Drug Name(s)

Morphine Sulfate Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
 - B. Patient has a diagnosis of pain due to sickle cell disease OR
 - C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. ALL of the following:
 - a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - d. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR
 - 3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND
 - e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND
 - f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Opioids ER PA - Oxycodone

Drug Name(s)

Oxycontin

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
 - B. Patient has a diagnosis of pain due to sickle cell disease OR
 - C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. ALL of the following:
 - a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - d. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR
 - 3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND
 - e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND
 - f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Opioids ER PA – Tramadol

Drug Name(s)

Tramadol Hcl Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
 - B. Patient has a diagnosis of pain due to sickle cell disease OR
 - C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. ALL of the following:
 - a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - d. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR
 - 3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND
 - e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND
 - f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oral Immunotherapy Agents PA – Oralair

Drug Name(s)

Oralair

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of allergic rhinitis, with or without conjunctivitis AND
- 2. Patient's diagnosis is confirmed with ONE of the following:
 - A. Positive skin test to ONE of the pollen extracts included in the requested agent OR
 - B. IgE specific antibodies to ONE of the extracts included in the requested agent: Sweet vernal, orchard, perennial rye, Timothy, or Kentucky blue grass AND
- 3. ONE of the following:
 - A. Patient has tried and had an inadequate response to TWO standard allergy medications, one of which was an intranasal corticosteroid OR
 - B. Patient has an intolerance or hypersensitivity to TWO standard allergy medications, one of which was an intranasal corticosteroid OR
 - C. Patient has an FDA labeled contraindication to TWO standard allergy medications, one of which was an intranasal corticosteroid AND
- 4. Patient will NOT be using the requested agent in combination with a subcutaneous injectable immunotherapy agent AND
- 5. The requested agent will be started, or has already been started, 3 to 4 months before the expected onset of the applicable pollen season AND
- 6. The first dose is given in the clinic/hospital under direct supervision from the provider for a period of at least 30 minutes AND
- 7. Patient has been prescribed epinephrine auto-injector for at home emergency use

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, immunologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Standard allergy medications:

Oral antihistamines, oral corticosteroids, intranasal corticosteroids, intranasal antihistamines, or leukotriene inhibitors

Orkambi PA

Drug Name(s)

Orkambi

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of cystic fibrosis AND
- 2. ONE of the following:
 - A. Patient has the presence of the F508del mutation on both alleles (homozygous) of the CFTR gene confirmed by genetic testing OR
 - B. Patient has another CFTR gene mutation(s) that is responsive to the requested agent, as indicated in the FDA label, confirmed by genetic testing AND
- 3. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of cystic fibrosis AND
- 3. Patient has had improvement or stabilization with the requested agent [e.g., improvement in FEV1 from baseline, increase in weight/BMI, improvement from baseline Cystic Fibrosis Questionnaire-Revised (CFQ-R) Respiratory Domain score, improvements in respiratory symptoms (cough, sputum production, and difficulty breathing), and/or reduced number of pulmonary exacerbations] AND
- 4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Otezla PA

Drug Name(s)

Otezla

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

- 1. BOTH of the following:
 - A. Patient has ONE of the following diagnoses:
 - i. Plaque psoriasis OR
 - ii. Active psoriatic arthritis AND
 - B. ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - iii. Patient's medication history indicates use of a biologic immunomodulator agent for the same FDA labeled indication OR
 - iv. Patient has tried and had an inadequate response to at least ONE conventional prerequisite agent for the requested indication OR
 - v. Patient has an intolerance or hypersensitivity to at least ONE conventional prerequisite agent for the requested indication OR
 - vi. Patient has an FDA labeled contraindication to at least ONE conventional prerequisite agent for the requested indication OR
- 2. Patient has a diagnosis of oral ulcers associated with Behcet's disease (BD)

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of plaque psoriasis, active psoriatic arthritis, or oral ulcers associated with Behcet's disease (BD) AND
- 3. Patient has had clinical benefit with the requested agent (slowing of disease progression or decrease in symptom severity and/or frequency)

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Formulary conventional agent required for diagnoses of plaque psoriasis or active psoriatic arthritis

Formulary conventional agents for plaque psoriasis include cyclosporine, methotrexate, tazarotene, topical calcitriol, or topical corticosteroids

Formulary conventional agents for active psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

NO prerequisites are required for a diagnosis of oral ulcers associated with Behcet's disease (BD)

Ozempic PA

Drug Name(s)

Ozempic

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of type 2 diabetes mellitus AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 - 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 5. BOTH of the following:
 - a. Patient has a diagnosis of established cardiovascular disease [e.g., myocardial infarction, stroke, any revascularization procedure, transient ischemic attack, unstable angina, amputation, symptomatic or asymptomatic coronary artery disease] AND
 - b. The requested agent will be used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Palynziq PA

Drug Name(s)

Palynziq

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of phenylketonuria (PKU) AND
- 2. Patient has a baseline blood Phe level greater than 600 micromol/L (10 mg/dL) AND
- 3. Patient will NOT be using the requested agent in combination with sapropterin for the requested indication AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of phenylketonuria (PKU) AND
- 3. ONE of the following:
 - a. Patient's blood Phe levels are being maintained within the acceptable range OR
 - b. Patient has had a decrease in blood Phe level from baseline AND
- 4. Patient will NOT be using the requested agent in combination with sapropterin for the requested indication AND
- 5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., metabolic or genetic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Initial approval will be for 9 months, renewal approval will be for 12 months

Panretin PA

Drug Name(s)

Panretin

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of cutaneous lesions associated with AIDS-related Kaposi's sarcoma (KS) OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. ONE of the following:
 - 1. BOTH of the following:
 - a. Patient has a diagnosis of cutaneous lesions associated with AIDS-related Kaposi's sarcoma (KS) AND
 - b. Patient does NOT require systemic anti-Kaposi's sarcoma therapy OR
 - 2. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, dermatologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Pegylated Interferon PA

Drug Name(s)

Pegasys

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic hepatitis B AND BOTH of the following:
 - i. The chronic hepatitis B infection has been confirmed by serological markers AND
 - ii. Patient has NOT been administered the requested agent for more than 48 weeks for the treatment of chronic hepatitis B OR
 - B. BOTH of the following:
 - i. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
 - ii. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype OR
 - C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

12 months for all other diagnoses. For hep B, hep C see Other Criteria

Other Criteria:

No prior peginterferon alfa use, approve 48 weeks for hepatitis B infection. Prior peginterferon alfa use, approve remainder of 48 weeks of total therapy for hepatitis B infection

Duration of therapy for hepatitis C: Based on FDA approved labeling or AASLD/IDSA guideline supported

Pirfenidone PA

Drug Name(s)

Pirfenidone

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
- 2. Patient has no known explanation for interstitial lung disease (ILD) or pulmonary fibrosis (e.g., radiation, drugs, metal dusts, sarcoidosis, or any connective tissue disease known to cause ILD)

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
- 3. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., pathologist, pulmonologist, radiologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Posaconazole PA

Drug Name(s)

Noxafil Pak

Posaconazole Inj

Posaconazole Dr

Posaconazole Susp

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of oropharyngeal candidiasis AND ONE of the following:
 - i. Patient has tried and had an inadequate response to fluconazole or an alternative antifungal agent OR
 - ii. Patient has an intolerance or hypersensitivity to fluconazole or an alternative antifungal agent OR
 - iii. Patient has an FDA labeled contraindication to fluconazole or an alternative antifungal agent OR
 - B. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida AND patient is severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
 - C. Patient has a diagnosis of invasive Aspergillus AND ONE of the following:
 - i. Patient has tried and had an inadequate response to an alternative antifungal agent OR
 - ii. Patient has an intolerance or hypersensitivity to an alternative antifungal agent OR
 - iii. Patient has an FDA labeled contraindication to an alternative antifungal agent OR
 - D. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

One month for oropharyngeal candidiasis, 6 months for all other indications

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

- A. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida and patient continues to be severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
- B. Patient has a diagnosis of invasive Aspergillus AND patient has continued indicators of active disease (e.g., continued radiologic findings, positive cultures, positive serum galactomannan assay for Aspergillus) OR
- C. BOTH of the following:
 - i. Patient has a diagnosis of oropharyngeal candidiasis AND
 - ii. Patient has had clinical benefit with the requested agent OR
- D. BOTH of the following:
 - i. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient has had clinical benefit with the requested agent

Prolia PA

Drug Name(s)

Prolia

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Osteopenia (osteoporosis prophylaxis)

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of:

1. ONE of:

A. Patient's (pt) sex is male or the pt is postmenopausal with a diagnosis of osteoporosis AND BOTH of:

- i. Pt's diagnosis was confirmed by ONE of:
 - 1. A fragility fracture in the hip or spine OR
 - 2. A T-score of -2.5 or lower OR
 - 3. A T-score of -1.0 to -2.5 AND ONE of:
 - a. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR
 - b. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR
 - c. A FRAX 10-year probability of hip fracture of 3% or greater AND

ii. ONE of:

- 1. Pt is at a very high fracture risk as defined by ONE of:
 - a. Pt had a recent fracture (within the past 12 months) OR
 - b. Pt had fractures while on FDA approved osteoporosis therapy OR
 - c. Pt has had multiple fractures OR
 - d. Pt had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR
 - e. Pt has a very low T-score (less than -3.0) OR
 - f. Pt is at high risk for falls or has a history of injurious falls OR
 - g. Pt has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR
- 2. ONE of:
 - a. Pt's medication history includes use of a bisphosphonate OR
 - b. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR
- B. Pt is requesting the agent for osteopenia (osteoporosis prophylaxis) AND ALL of:
 - i. ONE of:
 - 1. Pt's sex is male and the pt is 50 years of age or over OR
 - 2. Pt is postmenopausal AND

- ii. Pt has a T-score between -1.0 to -2.50 AND
- iii. ONE of:
 - a. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR
 - b. 10-year probability of a hip fracture 3% and greater per FRAX OR
 - c. 10-year probability of a major OP-related fracture 20% and greater per FRAX AND
- iv. ONE of:
 - a. Pt's medication history includes use of a bisphosphonate OR

Criteria continues: See Other Criteria

Age Restriction:

Prescriber Restrictions: Coverage Duration:

Approval will be for 12 months

- b. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR
- C. Pt's sex is a female with a diagnosis of breast cancer who is receiving aromatase inhibitor therapy AND ONE of:
 - i. Pt's medication history includes use of a bisphosphonate OR
 - ii. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR
- D. Pt's sex is male with a diagnosis of prostate cancer receiving androgen deprivation therapy (ADT) AND ONE of:
 - i. Pt's medication history includes use of a bisphosphonate OR
 - ii. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR
- E. Pt has a diagnosis of glucocorticoid-induced osteoporosis AND ALL of:
 - i. Pt is either initiating or continuing systemic glucocorticoids in a daily dose equivalent to 7.5 mg or greater of prednisone AND
 - ii. Pt is expected to remain on glucocorticoids for at least 6 months AND
 - iii. Pt's diagnosis was confirmed by ONE of:
 - 1. A fragility fracture in the hip or spine OR
 - 2. A T-score of -2.5 or lower OR
 - 3. A T-score of -1.0 to -2.5 AND ONE of the following:
 - a. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR
 - b. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR
 - c. A FRAX 10-year probability of hip fracture of 3% or greater AND
 - iv. ONE of:
 - 1. Pt is at a very high fracture risk as defined by ONE of the following:
 - a. Pt had a recent fracture (within the past 12 months) OR
 - b. Pt had fractures while on FDA approved osteoporosis therapy OR

- c. Pt has had multiple fractures OR
- d. Pt had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR
- e. Pt has a very low T-score (less than -3.0) OR
- f. Pt is at high risk for falls or has a history of injurious falls OR
- g. Pt has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR

2. ONE of:

- a. Pt's medication history includes use of a bisphosphonate OR
- b. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate AND

2. ONE of:

- A. Pt has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
- B. Pt has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
- C. Prescriber has indicated that the pt is not at risk for hypocalcemia (not including risk associated with the requested agent) AND
- 3. Pt will NOT be using the requested agent in combination with a bisphosphonate, another form of denosumab (e.g., Xgeva), romosozumab-aqqg, or parathyroid hormone analog (e.g., abaloparatide, teriparatide) for the requested indication AND
- 4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Promacta PA

Drug Name(s)

Promacta

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

- 1. Patient (pt) has a diagnosis of persistent or chronic immune (idiopathic) thrombocytopenia (ITP) AND ONE of the following:
 - A. Pt has tried and had an insufficient response to a corticosteroid or immunoglobulin (IVIg or anti-D) OR
 - B. Pt has an intolerance or hypersensitivity to a corticosteroid or immunoglobulin (IVIg or anti-D) OR
 - C. Pt has an FDA labeled contraindication to a corticosteroid or immunoglobulin (IVIg or anti-D) OR
 - D. Pt has had an insufficient response to a splenectomy OR
- 2. Pt has a diagnosis of hepatitis C associated thrombocytopenia AND ONE of the following:
 - A. Pt's platelet count is less than 75 x $10^9/L$ AND the intent is to increase platelet counts sufficiently to initiate pegylated interferon therapy OR
 - B. Pt is on concurrent therapy with a pegylated interferon and ribavirin AND is at risk for discontinuing hepatitis C therapy due to thrombocytopenia OR
- 3. Pt has a diagnosis of severe aplastic anemia (SAA) AND ALL of the following:
 - A. Pt has at least 2 of the following blood criteria:
 - i. Neutrophils less than 0.5 X 10^9/L OR
 - ii. Platelets less than 30 X 10^9/L OR
 - iii. Reticulocyte count less than 60 X 10^9/L AND
 - B. Pt has at least 1 of the following marrow criteria:
 - i. Severe hypocellularity is less than 25% OR
 - ii. Moderate hypocellularity is 25-50% with hematopoietic cells representing less than 30% of residual cells AND
 - C. ONE of the following:
 - i. Pt has tried and had an insufficient response to BOTH antithymocyte globulin (ATG) AND cyclosporine therapy OR
 - ii. BOTH of the following:
 - 1. Pt will be using the requested agent as first-line treatment (i.e., has not been treated with ATG and/or cyclosporine) AND
 - 2. Pt will use the requested agent in combination with standard immunosuppressive therapy (i.e., ATG AND cyclosporine) OR
- 4. Pt has another indication that is supported in CMS approved compendia for the requested agent **Age Restriction:**

Prescriber Restrictions:

Coverage Duration:

Initial: 6 months for ITP. Renewal: 12 months for ITP. Other indications, see Other Criteria.

Criteria for renewal approval require BOTH of the following:

- 1. Pt has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Pt has a diagnosis of persistent or chronic immune (idiopathic) thrombocytopenia (ITP) AND ONE of the following:
 - i. Pt's platelet count is 50 x 10^9/L or greater OR
 - ii. Pt's platelet count has increased sufficiently to avoid clinically significant bleeding OR
 - B. Pt has a diagnosis of hepatitis C associated thrombocytopenia AND BOTH of the following:
 - i. ONE of the following:
 - 1. Pt will be initiating hepatitis C therapy with pegylated interferon and ribavirin OR
 - 2. Pt will be maintaining hepatitis C therapy with pegylated interferon and ribavirin at the same time as the requested agent AND
 - ii. ONE of the following:
 - 1. Pt's platelet count is 90 x 10^9/L or greater OR
 - 2. Pt's platelet count has increased sufficiently to initiate or maintain pegylated interferon based therapy for the treatment of hepatitis C OR
 - C. Pt has a diagnosis of severe aplastic anemia (SAA) AND the pt has had clinical benefit with the requested agent OR
 - D. Pt has another indication that is supported in CMS approved compendia and the pt has had clinical benefit with the requested agent

Initial: 48 weeks for hepatitis C associated thrombocytopenia, 6 months for first-line therapy in severe aplastic anemia, 16 weeks for SAA, 12 months for All other indications

Renewal: 48 weeks for hepatitis C associated thrombocytopenia, 12 months for SAA, 12 months for All other indications

Pulmonary Hypertension PA – Adempas

Drug Name(s)

Adempas

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND
 - ii. Patient has an FDA labeled indication for the requested agent OR
 - B. Patient has a diagnosis of chronic thromboembolic pulmonary hypertension (CTEPH), WHO Group 4, as determined by a ventilation-perfusion scan and a confirmatory selective pulmonary angiography AND ALL of the following:
 - i. ONE of the following:
 - a. Patient is NOT a candidate for surgery OR
 - b. Patient has had pulmonary endarterectomy AND has persistent or recurrent disease AND
 - ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
 - iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
 - iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units OR C. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:
 - i. Patient's World Health Organization (WHO) functional class is II or greater AND
 - ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
 - iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
 - iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- v. ONE of the following:
 - a. The requested agent will be utilized as monotherapy OR
 - b. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), AND BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established pharmacotherapy AND
 - 2. The requested agent is in a different therapeutic class OR
 - c. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:
 - 1. ONE of the following:
 - i. A prostanoid has been started as one of the agents in the triple therapy OR
 - ii. Patient has an intolerance or hypersensitivity to a prostanoid OR
 - iii. Patient has an FDA labeled contraindication to a prostanoid AND
 - 2. Patient has unacceptable or deteriorating clinical status despite established pharmacotherapy AND
 - 3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Ambrisentan

Drug Name(s)

Ambrisentan

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND
 - ii. Patient has an FDA labeled indication for the requested agent OR
 - B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:
 - i. Patient's World Health Organization (WHO) functional class is II or greater AND
 - ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
 - iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
 - iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND
 - v. ONE of the following:
 - a. The requested agent will be utilized as monotherapy OR
 - b. The requested agent will be used in combination with a phosphodiesterase 5 (PDE5) inhibitor for dual therapy ONLY OR
 - c. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), [except for dual therapy requests for a phosphodiesterase 5 (PDE 5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:
 - 1. ONE of the following:
 - i. A prostanoid has been started as one of the agents in the triple therapy OR
 - ii. Patient has an intolerance or hypersensitivity to a prostanoid OR
 - iii. Patient has an FDA labeled contraindication to a prostanoid AND
 - 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Bosentan

Drug Name(s)

Bosentan

Tracleer

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Elevated liver enzymes accompanied by signs or symptoms of liver dysfunction/injury or a bilirubin level of 2 times the ULN (upper limit of normal) or greater AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1, as determined by right heart catheterization, AND ALL of the following:
 - i. Patient's World Health Organization (WHO) functional class is II or greater AND
 - ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
 - iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
 - iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND
 - v. ONE of the following:
 - a. The requested agent will be utilized as monotherapy OR
 - b. The requested agent will be used in combination with a phosphodiesterase 5 (PDE5) inhibitor for dual therapy ONLY OR
 - c. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), [except for dual therapy requests for a phosphodiesterase 5 inhibitor (PDE5) plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:
 - 1. ONE of the following:
 - i. A prostanoid has been started as one of the agents in the triple therapy OR
 - ii. Patient has an intolerance or hypersensitivity to a prostanoid OR
 - iii. Patient has an FDA labeled contraindication to a prostanoid AND
 - 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
- 3. All three agents in the triple therapy are from a different therapeutic class OR
- C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Opsumit

Drug Name(s)

Opsumit

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND
 - ii. Patient has an FDA labeled indication for the requested agent OR
 - B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:
 - i. Patient's World Health Organization (WHO) functional class is II or greater AND
 - ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
 - iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
 - iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND
 - v. ONE of the following:
 - a. The requested agent will be utilized as monotherapy OR
 - b. The requested agent will be used in combination with a phosphodiesterase 5 (PDE5) inhibitor for dual therapy ONLY OR
 - c. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), [except for dual therapy requests for a phosphodiesterase 5 (PDE 5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:
 - 1. ONE of the following:
 - i. A prostanoid has been started as one of the agents in the triple therapy OR
 - ii. Patient has an intolerance or hypersensitivity to a prostanoid OR
 - iii. Patient has an FDA labeled contraindication to a prostanoid
 - 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA - Sildenafil

Drug Name(s)

Sildenafil

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Concurrently taking another phosphodiesterase 5 (PDE 5) inhibitor with the requested agent AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:
 - i. Patient's World Health Organization (WHO) functional class is II or greater AND
 - ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
 - iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
 - iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND
 - v. ONE of the following:
 - a. The requested agent will be utilized as monotherapy OR
 - b. The requested agent will be used in combination with an endothelin receptor antagonist (ERA) for dual therapy ONLY OR
 - c. The requested agent will be utilized for add-on therapy to existing monotherapy, [except for dual requests for a phosphodiesterase 5 (PDE5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy) AND ALL of the following:
 - 1. ONE of the following:
 - i. A prostanoid has been started as one of the agents in the triple therapy OR
 - ii. Patient has an intolerance or hypersensitivity to a prostanoid OR
 - iii. Patient has an FDA labeled contraindication to a prostanoid AND
 - 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 3. All three agents in the triple therapy are from a different therapeutic class OR
- C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA - Tadalafil

Drug Name(s)

Alyq

Tadalafil 20Mg Tablet

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Concurrently taking another phosphodiesterase 5 (PDE 5) inhibitor with the requested agent AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:
 - i. Patient's World Health Organization (WHO) functional class is II or greater AND
 - ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
 - iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
 - iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND
 - v. ONE of the following:
 - a. The requested agent will be utilized as monotherapy OR
 - b. The requested agent will be used in combination with an endothelin receptor antagonist (ERA) for dual therapy ONLY OR
 - c. The requested agent will be utilized for add-on therapy to existing monotherapy, [except for dual requests for a phosphodiesterase 5 (PDE5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy) AND ALL of the following:
 - 1. ONE of the following:
 - i. A prostanoid has been started as one of the agents in the triple therapy OR
 - ii. Patient has an intolerance or hypersensitivity to a prostanoid OR
 - iii. Patient has an FDA labeled contraindication to a prostanoid AND
 - 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 3. All three agents in the triple therapy are from a different therapeutic class OR
- C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Ventavis

Drug Name(s)

Ventavis

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND
 - ii. Patient has an FDA labeled indication for the requested agent OR
 - B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:
 - i. Patient's World Health Organization (WHO) functional class is II or greater AND
 - ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
 - iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
 - iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND
 - v. ONE of the following:
 - a. The requested agent will be utilized as monotherapy OR
 - b. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), AND BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 2. The requested agent is in a different therapeutic class OR
 - c. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:
 - 1. Patient is WHO functional class III or IV AND
 - 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 3. All three agents in the triple therapy are from a different therapeutic class

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Drug is also subject to Part B versus Part D review.

Pyrimethamine PA

Drug Name(s)

Pyrimethamine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 6 months

Quinine PA

Drug Name(s)

Quinine Sulfate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

- 1. Patient has ONE of the following diagnoses:
 - a. Uncomplicated malaria OR
 - b. Babesiosis OR
 - c. An indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

7 days for malaria, 10 days for babesiosis, 12 months for all other diagnoses

Regranex PA

Drug Name(s)

Regranex

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - a. BOTH of the following:
 - i. Patient has a diagnosis of lower extremity diabetic neuropathic ulcer(s) that extends into the subcutaneous tissue or beyond AND
 - ii. The ulcer(s) intended for treatment has an adequate blood supply OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Repatha PA

Drug Name(s)

Repatha

Repatha Pushtronex System

Repatha Sureclick

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has ONE of the following:
 - A. A diagnosis of heterozygous familial hypercholesterolemia (HeFH) AND ONE of the following:
 - i. Genetic confirmation of one mutant allele at the LDLR, Apo-B, PCSK9, or 1/LDLRAP1 gene OR
 - ii. History of LDL-C greater than 190 mg/dL (greater than 4.9 mmol/L) (pretreatment) OR
 - iii. Patient has clinical manifestations of HeFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthoma, or xanthelasma) OR
 - iv. Patient has "definite" or "possible" familial hypercholesterolemia as defined by the Simon Broome criteria OR
 - v. Patient has a Dutch Lipid Clinic Network criteria score of greater than 5 OR
 - vi. Patient has a treated low-density lipoprotein cholesterol (LDL-C) level 100 mg/dL or greater after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy OR
 - B. A diagnosis of homozygous familial hypercholesterolemia (HoFH) AND ONE of the following:
 - i. Genetic confirmation of two mutant alleles at the LDLR, Apo-B, PCSK9, or LDLRAP1 gene OR
 - ii. History of untreated LDL-C greater than 500 mg/dL (greater than 13 mmol/L) or treated LDL-C 300 mg/dL or greater (7.76 mmol/L or greater) OR
 - iii. Patient has clinical manifestations of HoFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas, or xanthelasma) OR
 - C. A diagnosis of established cardiovascular disease [angina pectoris, coronary heart disease, myocardial infarction, transient ischemic attacks, cerebrovascular disease (CeVD) or peripheral vascular disease (PVD) or after coronary revascularization or carotid endarterectomy] AND the requested agent will be used to reduce the risk of myocardial infarction, stroke, and coronary revascularization OR
 - D. A diagnosis of primary hyperlipidemia (not associated with HeFH, HoFH, or established cardiovascular disease) OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

The agent was prescribed by, or in consultation with, a cardiologist, an endocrinologist, and/or a physician who focuses in the treatment of cardiovascular (CV) risk management and/or lipid disorders

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- E. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:
 - A. Patient has tried and had an inadequate response to a high-intensity statin (i.e., rosuvastatin 20-40 mg or atorvastatin 40-80 mg) OR
 - B. Patient has an intolerance to TWO different statins OR
 - C. Patient has an FDA labeled contraindication to a statin AND
- 4. Patient will NOT be using the requested agent in combination with another PCSK9 agent

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another PCSK9 agent

Ritalin PA

Drug Name(s)

Methylphenidate Hydrochloride (Ritalin)

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Roflumilast PA

Drug Name(s)

Roflumilast

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:
 - A. Patient has tried and had an inadequate response to an agent from two of the following categories:
 - i. long-acting beta-2 agonist (LABA) [e.g., salmeterol]
 - ii. long-acting muscarinic antagonist (LAMA) [e.g., umeclidinium]
 - iii. inhaled corticosteroid (ICS) [e.g., fluticasone] OR
 - B. Patient has an intolerance or hypersensitivity to an agent from two of the following categories:
 - i. long-acting beta-2 agonist (LABA) [e.g., salmeterol]
 - ii. long-acting muscarinic antagonist (LAMA) [e.g., umeclidinium]
 - iii. inhaled corticosteroid (ICS) [e.g., fluticasone] OR
 - C. Patient has an FDA labeled contraindication to an agent from two of the following categories:
 - i. long-acting beta-2 agonist (LABA) [e.g., salmeterol]
 - ii. long-acting muscarinic antagonist (LAMA) [e.g., umeclidinium]
 - iii. inhaled corticosteroid (ICS) [e.g., fluticasone]

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Rybelsus PA

Drug Name(s)

Rybelsus

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of type 2 diabetes mellitus AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 - 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Samsca PA

Drug Name(s)

Tolvaptan

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the request agent AND Any underlying liver disease, including cirrhosis **Required Medical Information:**

Criteria for approval require ALL of the following:

- 1. The requested agent was initiated (or re-initiated) in the hospital AND
- 2. Prior to initiating the requested agent, the patient has or had a diagnosis of clinically significant hypervolemic or euvolemic hyponatremia defined by ONE of the following:
 - A. Serum sodium is less than 125 mEq/L OR
 - B. Serum sodium is 125 mEq/L or greater AND patient has symptomatic hyponatremia that has resisted correction with fluid restriction AND
- 3. Medications known to cause hyponatremia have been evaluated and discontinued when appropriate AND
- 4. Patient has NOT already received 30 days of therapy with the requested agent following the most recent hospitalization for initiation of therapy AND
- 5. The requested dose is within the FDA labeled dosing for the requested indication (Recommended starting dose is 15 mg once daily. Dosage may be increased at intervals greater than or equal to 24 hours to 30 mg once daily, and to a maximum of 60 mg once daily as needed to raise serum sodium. Do not administer for more than 30 days to minimize the risk of liver injury.)

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 30 days

Sapropterin PA

Drug Name(s)

Javygtor

Sapropterin Dihydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of phenylketonuria (PKU) AND
- 2. Prescriber has submitted a baseline blood Phe level measured prior to initiation of therapy with the requested agent, which is above the recommended levels indicated for the patient's age range or condition AND
- 3. Patient will NOT be using the requested agent in combination with Palynziq (pegvaliase-pqpz) for the requested indication AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of phenylketonuria (PKU) AND
- 3. ONE of the following:
 - a. Patient's blood Phe levels are being maintained within the acceptable range OR
 - b. Patient has had a decrease in blood Phe level from baseline AND
- 4. Patient will NOT be using the requested agent in combination with Palynziq (pegvaliase-pqpz) for the requested indication AND
- 5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., metabolic or genetic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Initial: 2 months if dose is 5 to less than 20 mg/kg/day, 1 month if 20 mg/kg/day Renewal: 12 months Other Criteria:

Self - Administered Oncology PA

Drug Name(s)

Abiraterone Acetate

Akeega

Alecensa

Alunbrig

Augtyro

Ayvakit

Balversa

Besremi

Bexarotene Cap

Bosulif

Braftovi

Brukinsa

Cabometyx

Calquence

Caprelsa

Cometriq

Copiktra

Cotellic

Daurismo

Erivedge

Erleada

Erlotinib Hydrochloride

Everolimus

Exkivity

Farydak

Fotivda

Fruzagla

Gavreto

Gefitinib

Gilotrif

Ibrance

Iclusig

Idhifa

Imatinib Mesylate

Imbruvica

Inlyta

Inqovi

Inrebic

Iwilfin

Jakafi

Jaypirca

Kisqali

Kisqali Femara 200 Dose

Kisqali Femara 400 Dose

Kisqali Femara 600 Dose

Koselugo

Krazati

Lapatinib Ditosylate

Lenalidomide

Lenvima 10 Mg Daily Dose

Lenvima 12Mg Daily Dose

Lenvima 14 Mg Daily Dose

Lenvima 18 Mg Daily Dose

Lenvima 20 Mg Daily Dose

Lenvima 24 Mg Daily Dose

Lenvima 4 Mg Daily Dose

Lenvima 8 Mg Daily Dose

Lonsurf

Lorbrena

Lumakras

Lynparza

Lytgobi

Matulane

Mekinist

Mektovi

Nerlynx

Ninlaro

Nubeqa

Odomzo

Ogsiveo

Ojjaara

Ojjuuru

Onureg

Orgovyx

Orserdu

Pazopanib

Pemazyre

Piqray 200Mg Daily Dose

Piqray 250Mg Daily Dose

Pigray 300Mg Daily Dose

Pomalyst

Qinlock

Retevmo

Revlimid

Rezlidhia

Rozlytrek

Rubraca

Rydapt

Scemblix

Sorafenib

Sprycel

Stivarga

Sunitinib Malate

Tabrecta

Tafinlar

Tagrisso

Talzenna

Tasigna

Tazverik

Tepmetko

Thalomid

Tibsovo

Tretinoin 10Mg Cap

Truqap

Tukysa

Turalio

Vanflyta

Venclexta

Venclexta Starting Pack

Verzenio

Vitrakvi

Vizimpro

Vonjo

Votrient

Welireg

Xalkori

Xospata

Xpovio

Xpovio 100 Mg Once Weekly

Xpovio 40 Mg Once Weekly

Xpovio 40 Mg Twice Weekly

Xpovio 60 Mg Once Weekly

Xpovio 60 Mg Twice Weekly

Xpovio 80 Mg Once Weekly

Xpovio 80 Mg Twice Weekly

Xtandi

Yonsa

Zejula

Zelboraf

Zolinza

Zydelig Zykadia

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent AND iii. ONE of the following:
 - a. The requested agent is FDA labeled or supported by compendia as a first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or compendia supported therapy that are indicated as first-line therapy for the requested indication OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iv. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines AND

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

- v. ONE of the following:
 - a. The requested agent is not Bosulif or Tasigna OR
 - b. The requested agent is Bosulif or Tasigna AND ONE of the following:
 - 1. Patient's medication history indicates use of imatinib OR Sprycel for the requested indication (if applicable) OR
 - 2. Patient has an intolerance or hypersensitivity to imatinib OR Sprycel OR

- 3. Patient has an FDA labeled contraindication to imatinib OR Sprycel OR
- 4. CMS approved compendia does not support the use of imatinib OR Sprycel for the requested indication OR
- 5. Prescriber has provided information in support of use of Bosulif or Tasigna over imatinib OR Sprycel for the requested indication AND
- vi. ONE of the following:
 - a. The requested agent is not Calquence OR
 - b. The requested agent is Calquence AND ONE of the following:
 - 1. Patient's medication history indicates use of Brukinsa OR Imbruvica for the requested indication (if applicable) OR
 - 2. Patient has an intolerance or hypersensitivity to Brukinsa OR Imbruvica OR
 - 3. Patient has an FDA labeled contraindication to Brukinsa OR Imbruvica
 - 4. CMS approved compendia do not support the use of Brukinsa OR Imbruvica for the requested indication OR
 - 5. Prescriber has provided information in support of use of Calquence over Brukinsa OR Imbruvica for the requested indication

Signifor LAR PA

Drug Name(s)

Signifor Lar

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Severe hepatic impairment (i.e., Child Pugh C)

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of acromegaly AND ONE of the following:
 - i. Patient had an inadequate response to surgery as indicated by growth hormone and serum IGF-1 levels that are above the reference ranges for the patient's gender and age OR
 - ii. Patient is NOT a candidate for surgery OR
 - B. Patient has a diagnosis of Cushing's disease (CD) AND ONE of the following:
 - i. Patient had an inadequate response to pituitary surgical resection OR
 - ii. Patient is NOT a candidate for pituitary surgical resection OR
 - C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of acromegaly AND ONE of the following:
 - i. Patient has growth hormone and serum IGF-1 levels that are within normal limits for patient's gender and age reference range OR
 - ii. Patient has had clinical improvement (e.g., reduction in tumor size, decreased headaches, improved cardiovascular or respiratory symptoms) OR
 - B. Patient has a diagnosis of Cushing's disease (CD) AND BOTH of the following:
 - i. Patient has a urinary free cortisol level less than or equal to the upper limit of normal AND
 - ii. Patient has had improvement in at least ONE of the following clinical signs and symptoms:
 - 1. Fasting plasma glucose OR
 - 2. Hemoglobin A1c OR
 - 3. Hypertension OR
 - 4. Weight OR
 - C. BOTH of the following:
 - i. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Initial: Acromegaly - 6 months, CD - 7 months, All other diagnoses - 12 months, Renewal: 12 months

Signifor PA

Drug Name(s)

Signifor

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Severe hepatic impairment (i.e., Child Pugh C)

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of Cushing's disease (CD) AND ONE of the following:
 - i. Patient had an inadequate response to pituitary surgical resection OR
 - ii. Patient is NOT a candidate for pituitary surgical resection OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of Cushing's disease (CD) AND BOTH of the following:
 - i. Patient has a urinary free cortisol level less than or equal to the upper limit of normal AND
 - ii. Patient has had improvement in at least ONE of the following clinical signs and symptoms:
 - 1. Fasting plasma glucose OR
 - 2. Hemoglobin A1c OR
 - 3. Hypertension OR
 - 4. Weight OR
 - B. BOTH of the following:
 - i. Patient has an indication that is supported in CMS approved compendia for the requested agent $\ensuremath{\mathsf{AND}}$
 - ii. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Initial approval: 6 months for CD, 12 months for all other diagnoses, Renewal approval: 12 months Other Criteria:

Sivextro PA

Drug Name(s)

Sivextro

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following:
 - A. BOTH of the following:
 - i. A documented acute bacterial skin and skin structure infection (ABSSSI) defined as a bacterial infection of the skin with a lesion size area of at least 75 cm2 (lesion size measured by the area of redness, edema, or induration) AND
 - ii. The infection is due to Staphylococcus aureus, Streptococcus pyogenes, Streptococcus agalactiae, Streptococcus anginosus, Streptococcus intermedius, Streptococcus constellatus, or Enterococcus faecalis OR
 - B. Another indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. The requested agent is prescribed by an infectious disease specialist or the prescriber has consulted with an infectious disease specialist on treatment of this patient OR
 - B. The requested agent is NOT prescribed by an infectious disease specialist or the prescriber has NOT consulted with an infectious disease specialist on treatment of this patient AND ONE of the following:
 - i. There is documentation of resistance to TWO of the following: beta-lactams, macrolides, clindamycin, tetracycline, or co-trimoxazole at the site of infection OR
 - ii. Patient has an intolerance or hypersensitivity to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
 - iii. Patient has an FDA labeled contraindication to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
 - iv. There is documentation of resistance to vancomycin at the site of infection OR
 - v. Patient has an intolerance or hypersensitivity to vancomycin OR
 - vi. Patient has an FDA labeled contraindication to vancomycin AND
- 3. Patient will NOT be using the requested agent in combination with linezolid for the same infection
- 4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 days for ABSSSI or 30 days for all other indications

Sodium Oxybate PA

Drug Name(s)

Sodium Oxybate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of narcolepsy with cataplexy OR
 - B. Patient has a diagnosis of narcolepsy with excessive daytime sleepiness AND BOTH of the following:
 - i. ONE of the following:
 - a. Patient is under 18 years of age OR
 - b. ONE of the following:
 - 1. Patient has tried and had an inadequate response to modafinil or armodafinil OR
 - 2. Patient has an intolerance or hypersensitivity to modafinil or armodafinil OR
 - 3. Patient has an FDA labeled contraindication to modafinil or armodafinil AND
 - ii. ONE of the following:
 - a. Patient has tried and had an inadequate response to ONE standard stimulant agent (e.g., methylphenidate) OR
 - b. Patient has an intolerance or hypersensitivity to ONE standard stimulant agent (e.g., methylphenidate) OR
 - c. Patient has an FDA labeled contraindication to ONE standard stimulant agent (e.g., methylphenidate) OR
 - C. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Patient is 7 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Somatostatin Analogs PA - Lanreotide

Drug Name(s)

Somatuline Depot

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. ONE of the following:
 - A. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent OR
 - B. ONE of the following:
 - i. Patient has a diagnosis of acromegaly AND ONE of the following:
 - a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR
 - b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR
 - c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by growth hormone levels or serum IGF-1 levels that are above the reference range OR
 - ii. Patient has a diagnosis of gastroenteropancreatic neuroendocrine tumors AND BOTH of the following:
 - a. The tumors are well or moderately differentiated AND
 - b. ONE of the following:
 - 1. The tumors are unresectable, locally advanced OR
 - 2. Patient has metastatic disease OR
 - iii. Patient has a diagnosis of carcinoid syndrome OR
 - iv. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
- 2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - 1. Patient has a diagnosis of acromegaly OR
 - 2. Patient has a diagnosis of metastatic OR unresectable, locally advanced, well or moderately differentiated gastroenteropancreatic neuroendocrine tumors OR
 - 3. Patient has a diagnosis of carcinoid syndrome OR
 - 4. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient has had clinical benefit with the requested agent AND
- 4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Somatostatin Analogs PA – Octreotide

Drug Name(s)

Octreotide Acetate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. ONE of the following:
 - A. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - B. ONE of the following:
 - i. Patient has a diagnosis of acromegaly AND ONE of the following:
 - a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR
 - b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR
 - c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by growth hormone levels or serum IGF-1 levels that are above the reference range OR
 - ii. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR
 - iii. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR
 - iv. Patient has a diagnosis of dumping syndrome AND ONE of the following:
 - a. Patient has tried and had an inadequate response to acarbose OR
 - b. Patient has an intolerance or hypersensitivity to acarbose OR
 - c. Patient has an FDA labeled contraindication to acarbose OR
 - v. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
- 2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of acromegaly OR
 - B. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR
 - C. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR
 - D. Patient has a diagnosis of dumping syndrome OR
 - E. Patient has another indication that is supported in CMS approved compendia for the requested agent

AND

- 3. Patient has had clinical benefit with the requested agent AND
- 4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Somatostatin Analogs PA - Somavert

Drug Name(s)

Somavert

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has a diagnosis of acromegaly AND ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. BOTH of the following:
 - i. ONE of the following:
 - a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR
 - b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR
 - c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by serum IGF-1 levels that are above the reference range AND
 - ii. ONE of the following:
 - a. Patient has tried and had an inadequate response to octreotide or Somatuline Depot (lanreotide) OR
 - b. Patient has an intolerance or hypersensitivity to octreotide or Somatuline Depot (lanreotide) OR
 - c. Patient has an FDA labeled contraindication to octreotide or Somatuline Depot (lanreotide) AND
- 2. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of acromegaly AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal

Sovaldi PA

Drug Name(s)

Sovaldi

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND
- 2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
- 3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
- 4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND
- 5. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR
 - D. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported **Other Criteria**:

Spravato PA

Drug Name(s)

Spravato 56Mg Dose

Spravato 84Mg Dose

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. BOTH of the following:
 - a. ONE of the following:
 - i. Patient has a diagnosis of treatment-resistant depression (TRD) OR
 - ii. Patient has a diagnosis of major depressive disorder (MDD) with acute suicidal ideation or behavior AND
 - b. The requested agent will be used in combination with an oral antidepressant AND
- 2. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent OR
 - c. ALL of the following:
 - i. ONE of the following:
 - 1. BOTH of the following:
 - a. Patient has a diagnosis of treatment-resistant depression (TRD) AND
 - b. Patient has tried and had an inadequate response to at least two different oral antidepressants (e.g., SSRIs, SNRIs) OR
 - 2. Patient has a diagnosis of major depressive disorder (MDD) with acute suicidal ideation or behavior AND
 - ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., psychiatrist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Strensiq PA

Drug Name(s)

Strensiq

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Perinatal or infantile-onset hypophosphatasia OR
 - B. Juvenile-onset hypophosphatasia AND
- 2. Patient has documentation (i.e., medical records) of clinical manifestations to support the diagnosis of hypophosphatasia at the age of onset prior to age 18 (e.g., vitamin B6-dependent seizures, skeletal abnormalities such as rachitic chest deformity leading to respiratory problems or bowed arms/legs, "failure to thrive") AND
- 3. Patient has documentation (i.e., medical records) of radiographic imaging to support the diagnosis of hypophosphatasia at the age of onset prior to age 18 (e.g., infantile rickets, alveolar bone loss, craniosynostosis, fractures) AND
- 4. Patient has documentation (i.e., medical records) of confirmed mutation(s) in the ALPL gene that encodes the tissue non-specific isoenzyme of alkaline phosphatase (TNSALP) AND
- 5. Patient has documentation (i.e., medical records) of a measured total serum alkaline phosphatase (ALP) level that is below the normal lab reference range for age and sex AND
- 6. Patient has documentation (i.e., medical records) of ONE of the following:
 - A. Elevated urine concentration of phosphoethanolamine (PEA) OR
 - B. Elevated serum concentration of pyridoxal 5'-phosphate (PLP) in the absence of vitamin supplements within one week prior to the test OR
 - C. Elevated urinary inorganic pyrophosphate (PPi) AND
- 7. The requested dose is within FDA labeled dosing (based on the patient's weight) for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist or geneticist with expertise in metabolic bone diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has ONE of the following diagnoses:
 - A. Perinatal or infantile-onset hypophosphatasia OR

- B. Juvenile-onset hypophosphatasia AND
- 3. There is documentation (i.e., medical records) that the patient has had a decrease from baseline (before treatment with the requested agent) in at least ONE of the following levels:
 - a. Urine concentration of phosphoethanolamine (PEA) OR
 - b. Serum concentration of pyridoxal 5'-phosphate (PLP) in the absence of vitamin supplements within one week prior to the test OR
 - c. Urinary inorganic pyrophosphate (PPi) AND
- 4. Patient has documentation (i.e., medical records) of clinical improvement and/or stabilization with the requested agent (e.g., improvement in respiratory status, growth, pain, radiographic findings, other symptoms associated with the disease) AND
- 5. The requested dose is within FDA labeled dosing (based on the patient's weight) for the requested indication

Substrate Reduction Therapy PA - Miglustat

Drug Name(s)

Miglustat

Yargesa

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of Gaucher disease type 1 (GD1) confirmed by ONE of the following:
 - A. A baseline glucocerebrosidase enzyme activity of less than or equal to 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR
 - B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND
- 2. Prescriber has drawn baseline measurements of hemoglobin level, platelet count, liver volume, and spleen volume AND
- 3. Prior to any treatment for the intended diagnosis, the patient has had at least ONE of the following clinical presentations:
 - A. Anemia [defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender] OR
 - B. Thrombocytopenia (defined as platelet count of less than 100,000 per microliter) OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth failure (i.e., growth velocity is below the standard mean for age) OR
 - F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of Gaucher disease type 1 (GD1) AND
- 3. Patient has had improvements or stabilization with the requested agent as indicated by ONE of the following:
 - A. Spleen volume OR
 - B. Hemoglobin level OR
 - C. Liver volume OR
 - D. Platelet count OR
 - E. Growth OR
 - F. Bone pain or crisis

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Tasimelteon Capsule PA

Drug Name(s)

Tasimelteon

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. Patient has a diagnosis of Non-24-hour sleep-wake disorder AND
 - ii. Patient is totally blind (i.e., no light perception) OR
 - B. BOTH of the following:
 - i. Patient has a diagnosis of Smith-Magenis Syndrome (SMS) confirmed by the presence of ONE of the following genetic mutations:
 - A. A heterozygous deletion of 17p11.2 OR
 - B. A heterozygous pathogenic variant involving RAI1 AND
 - ii. The requested agent is being used to treat nighttime sleep disturbances associated with SMS

Age Restriction:

For diagnosis of Non-24-hour sleep-wake disorder, patient is 18 years of age or over. For diagnosis of Smith-Magenis Syndrome (SMS), patient is 16 years of age or over.

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist, sleep specialist, psychiatrist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Teriparatide PA

Drug Name(s)

Forteo

Teriparatide

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Postmenopausal with osteoporosis OR
 - B. Patient's sex is male with primary or hypogonadal osteoporosis OR
 - C. Osteoporosis with sustained systemic glucocorticoid therapy AND
- 2. Patient's diagnosis was confirmed by ONE of the following:
 - A. A fragility fracture in the hip or spine OR
 - B. A T-score of -2.5 or lower OR
 - C. A T-score of -1.0 to -2.5 AND ONE of the following:
 - i. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR
 - ii. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR
 - iii. A FRAX 10-year probability of hip fracture of 3% or greater AND
- 3. ONE of the following:
 - A. Patient is at a very high fracture risk as defined by ONE of the following:
 - i. Patient had a recent fracture (within the past 12 months) OR
 - ii. Patient had fractures while on FDA approved osteoporosis therapy OR
 - iii. Patient has had multiple fractures OR
 - iv. Patient had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR
 - v. Patient has a very low T-score (less than -3.0) OR
 - vi. Patient is at high risk for falls or has a history of injurious falls OR
 - vii. Patient has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR
 - B. ONE of the following:
 - i. Patient has tried and had an inadequate response to a bisphosphonate OR
 - ii. Patient has an intolerance or hypersensitivity to a bisphosphonate OR
 - iii. Patient has an FDA labeled contraindication to a bisphosphonate AND
- 4. Patient will NOT be using the requested agent in combination with a bisphosphonate, denosumab (e.g., Prolia, Xgeva), romosozumab-aqqg, or another parathyroid hormone analog (e.g., abaloparatide) for the requested indication AND

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

No prior teriparatide and/or Tymlos use approve 2 years, Prior use - see Other Criteria Other Criteria:

- 5. The requested dose is within FDA labeled dosing for the requested indication AND $\,$
- 6. ONE of the following:
 - A. Patient has never received treatment with teriparatide or Tymlos (abaloparatide) OR
 - B. Patient has been previously treated with teriparatide or Tymlos (abaloparatide) AND ONE of the following:
 - i. The total cumulative duration of treatment with teriparatide and Tymlos (abaloparatide) has NOT exceeded 2 years OR
 - ii. Patient has received 2 years or more of treatment with teriparatide, and remains at or has returned to having a high risk for fracture

Prior teriparatide and/or Tymlos use approve remainder of 2 years of total cumulative therapy. Approve 1 year if patient has received 2 years or more teriparatide

Tetrabenazine PA

Drug Name(s)

Tetrabenazine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chorea associated with Huntington's disease OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. Patient does NOT have a current diagnosis of depression OR
 - B. Patient has a current diagnosis of depression and is being treated for depression AND
- 3. ONE of the following:
 - A. Patient does NOT have a diagnosis of suicidal ideation and/or behavior OR
 - B. Patient has a diagnosis of suicidal ideation and/or behavior and must NOT be actively suicidal AND
- 4. Patient will NOT be using the requested agent in combination with a monoamine oxidase inhibitor (MAOI) AND
- 5. Patient will NOT be using the requested agent in combination with reserpine

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Tobramycin neb PA

Drug Name(s)

Tobramycin Neb

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of cystic fibrosis AND
- 2. Documentation has been provided that indicates the patient has a Pseudomonas aeruginosa respiratory infection AND
- 3. ONE of the following:
 - a. Patient is NOT currently (within the past 60 days) being treated with another inhaled antibiotic (e.g., inhaled aztreonam, inhaled tobramycin) OR
 - b. Patient is currently (within the past 60 days) being treated with another inhaled antibiotic (e.g., inhaled aztreonam, inhaled tobramycin) AND ONE of the following:
 - i. Prescriber has confirmed that the other inhaled antibiotic will be discontinued, and that therapy will be continued only with the requested agent OR
 - ii. Prescriber has provided information in support of another inhaled antibiotic therapy used concurrently with or alternating with (i.e., continuous alternating therapy) the requested agent

Drug is also subject to Part B versus Part D review.

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Topical Diclofenac 3% Gel PA

Drug Name(s)

Diclofenac Sodium 3% Gel

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has a diagnosis of actinic keratosis (AK)

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 3 months

Topical Retinoids PA – Tazarotene

Drug Name(s)

Tazarotene

Tazorac

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for cosmetic purposes

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - a. Patient has an FDA labeled indication for the requested agent OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Topical Retinoids PA – Tretinoin

Drug Name(s)

Avita

Tretinoin Cream, Gel

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for cosmetic purposes

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - a. Patient has an FDA labeled indication for the requested agent OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Trelstar PA

Drug Name(s)

Trelstar Mixject

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. BOTH of the following:
 - i. Patient is NOT currently being treated with the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent AND
- 3. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Trientine PA

Drug Name(s)

Trientine Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has a diagnosis of Wilson's disease confirmed by ONE of the following:
 - A. Confirmation of genetic mutation of the ATP7B gene OR
 - B. Patient has TWO or more of the following:
 - i. Presence of hepatic abnormality (e.g., acute liver failure, cirrhosis, fatty liver)
 - ii. Presence of Kayser-Fleischer rings
 - iii. Serum ceruloplasmin level less than 20 mg/dL
 - iv. Basal urinary copper excretion greater than 40 mcg/24 hours or the testing laboratory's upper limit of normal
 - v. Hepatic parenchymal copper content greater than 40 mcg/g dry weight
 - vi. Presence of neurological symptoms (e.g., dystonia, hypertonia, rigidity with tremors, dysarthria, muscle spasms, dysphasia, polyneuropathy, dysautonomia) AND
- 2. ONE of the following:
 - A. Patient has tried and had an inadequate response to penicillamine OR
 - B. Patient has an intolerance or hypersensitivity to penicillamine OR
 - C. Patient has an FDA labeled contraindication to penicillamine

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of Wilson's disease AND
- 3. Patient has had clinical benefit with the requested agent as evidenced by ONE of the following:
 - A. Improvement and/or stabilization in hepatic abnormality OR
 - B. Reduction in Kayser-Fleischer rings OR
 - C. Improvement and/or stabilization in neurological symptoms (e.g., dystonia, hypertonia, rigidity with tremors, dysarthria, muscle spasms, dysphasia, polyneuropathy, dysautonomia) OR
 - D. Basal urinary copper excretion greater than 200 mcg/24 hours

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist, neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Trikafta PA

Drug Name(s)

Trikafta

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of cystic fibrosis AND
- 2. ONE of the following:
 - A. Patient has the presence of the F508del mutation in at least ONE allele (heterozygous OR homozygous) of the CFTR gene confirmed by genetic testing OR
 - B. Patient has ONE of the CFTR gene mutations or a mutation in the CFTR gene that is responsive based on in vitro data, as indicated in the FDA label, confirmed by genetic testing OR
 - C. Patient has another CFTR gene mutation(s) that is responsive to the requested agent, as indicated in the FDA label, confirmed by genetic testing AND
- 3. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of cystic fibrosis AND
- 3. Patient has had improvement or stabilization with the requested agent [e.g., improvement in FEV1 from baseline, increase in weight/BMI, improvement from baseline Cystic Fibrosis Questionnaire-Revised (CFQ-R) Respiratory Domain score, improvements in respiratory symptoms (cough, sputum production, and difficulty breathing), and/or reduced number of pulmonary exacerbations] AND
- 4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Trulicity PA

Drug Name(s)

Trulicity

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of type 2 diabetes mellitus AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 - 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 5. BOTH of the following:
 - a. Patient has a diagnosis of established cardiovascular disease [e.g., myocardial infarction, stroke, any revascularization procedure, transient ischemic attack, unstable angina, amputation, symptomatic or asymptomatic coronary artery disease] AND
 - b. The requested agent will be used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Tymlos PA

Drug Name(s)

Tymlos

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient (pt) has ONE of the following:
 - A. Postmenopausal with osteoporosis OR
 - B. Pt's sex is male with osteoporosis AND
- 2. BOTH of the following:
 - A. Pt's diagnosis was confirmed by ONE of the following:
 - i. A fragility fracture in the hip or spine OR
 - ii. A T-score of -2.5 or lower OR
 - iii. A T-score of -1.0 to -2.5 AND ONE of the following:
 - a. A fragility fracture of proximal humerus, pelvis, or distal forearm OR
 - b. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR
 - c. A FRAX 10-year probability of hip fracture of 3% or greater AND
 - B. ONE of the following:
 - i. Pt is at a very high fracture risk as defined by ONE of the following:
 - a. Pt had a recent fracture (within the past 12 months) OR
 - b. Pt had fractures while on FDA approved osteoporosis therapy OR
 - c. Pt has had multiple fractures OR
 - d. Pt had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR
 - e. Pt has a very low T-score (less than -3.0) OR
 - f. Pt is at high risk for falls or has a history of injurious falls OR
 - g. Pt has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR
 - ii. ONE of the following:
 - a. Pt has tried and had an inadequate response to a bisphosphonate OR
 - b. Pt has an intolerance or hypersensitivity to a bisphosphonate OR
 - c. Pt has an FDA labeled contraindication to a bisphosphonate AND
- 3. Pt will NOT be using the requested agent in combination with a bisphosphonate, denosumab (e.g., Prolia, Xgeva), romosozumab-aqqg, or another parathyroid hormone analog (e.g., teriparatide) for the requested indication AND
- 4. The requested dose is within FDA labeled dosing for the requested indication AND
- 5. The total cumulative duration of treatment with teriparatide and Tymlos (abaloparatide) has not exceeded 2 years

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

No prior Tymlos and/or teriparatide use approve 2 years, Prior use - see Other Criteria

Other Criteria:

Prior Tymlos and/or teriparatide use approve remainder of 2 years of total cumulative therapy

Ubrelvy PA

Drug Name(s)

Ubrelvy

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of migraine AND
- 2. The requested agent is being used for the treatment of acute migraine with or without aura AND
- 3. ONE of the following:
 - A. Patient has tried and had an inadequate response to a triptan (e.g., sumatriptan, rizatriptan) agent OR
 - B. Patient has an intolerance, or hypersensitivity to a triptan OR
 - C. Patient has an FDA labeled contraindication to a triptan AND
- 4. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP)

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of migraine AND
- 3. The requested agent is being used for the treatment of acute migraine with or without aura AND
- 4. Patient has had clinical benefit with the requested agent AND
- 5. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP)

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Urea Cycle Disorders PA - Sodium Phenylbutyrate

Drug Name(s)

Sodium Phenylbutyrate

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of ONE of the following:
- a. Urea cycle disorder with neonatal-onset involving deficiencies of carbamylphosphate synthetase, ornithine transcarbamylase, or argininosuccinic acid synthetase OR
- b. Urea cycle disorder with late-onset and history of hyperammonemic encephalopathy involving deficiencies of carbamylphosphate synthetase, ornithine transcarbamylase, or argininosuccinic acid synthetase AND
- 2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., geneticist, metabolic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Viberzi PA

Drug Name(s)

Viberzi

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has a diagnosis of irritable bowel syndrome with diarrhea (IBS-D)

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Victoza PA

Drug Name(s)

Victoza

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of type 2 diabetes mellitus AND
- 2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 - 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 - 5. BOTH of the following:
 - a. Patient has a diagnosis of established cardiovascular disease [e.g., myocardial infarction, stroke, any revascularization procedure, transient ischemic attack, unstable angina, amputation, symptomatic or asymptomatic coronary artery disease] AND
 - b. The requested agent will be used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Voriconazole PA

Drug Name(s)

Voriconazole

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of invasive Aspergillus OR
 - B. Patient has a serious infection caused by Scedosporium apiospermum or Fusarium species OR
 - C. Patient has a diagnosis of esophageal candidiasis or candidemia in nonneutropenic patient AND ONE of the following:
 - i. Patient has tried and had an inadequate response to fluconazole or an alternative antifungal agent OR
 - ii. Patient has an intolerance or hypersensitivity to fluconazole or an alternative antifungal agent OR
 - iii. Patient has an FDA labeled contraindication to fluconazole or an alternative antifungal agent OR
 - D. Patient has a diagnosis of blastomycosis AND ONE of the following:
 - i. Patient has tried and had an inadequate response to itraconazole OR
 - ii. Patient has an intolerance or hypersensitivity to itraconazole OR
 - iii. Patient has an FDA labeled contraindication to itraconazole OR
 - E. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida AND patient is severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
 - F. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

One month for esophageal candidiasis, 6 months for all other indications

Other Criteria:

Criteria for renewal approval require BOTH of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:

- A. Patient has a diagnosis of invasive Aspergillus, a serious infection caused by Scedosporium apiospermum or Fusarium species, esophageal candidiasis, candidemia in nonneutropenic patient, or blastomycosis and patient has continued indicators of active disease (e.g., continued radiologic findings, positive cultures, positive serum galactomannan assay for Aspergillus) OR B. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida and patient continues to be severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
- C. BOTH of the following:
 - i. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient has had clinical benefit with the requested agent

Vosevi PA

Drug Name(s)

Vosevi

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND
- 2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
- 3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
- 4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND
- 5. If genotype 1, the patient's subtype has been identified and provided

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Vyndamax PA

Drug Name(s)

Vyndamax

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
- 2. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
- 3. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
- 3. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
- 4. Patient has had clinical benefit with the requested agent AND
- 5. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Vyndaqel PA

Drug Name(s)

Vyndagel

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
- 2. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
- 3. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
- 3. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
- 4. Patient has had clinical benefit with the requested agent AND
- 5. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Xermelo PA

Drug Name(s)

Xermelo

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of carcinoid syndrome diarrhea AND
- 2. Patient has tried and had an inadequate response to treatment with a somatostatin analog (e.g., Sandostatin [octreotide], Sandostatin LAR [octreotide], Somatuline Depot [lanreotide]) AND
- 3. The requested agent will be used in combination with a somatostatin analog (e.g., Sandostatin [octreotide], Sandostatin LAR [octreotide], Somatuline Depot [lanreotide])

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of carcinoid syndrome diarrhea AND
- 3. Patient has had clinical benefit with the requested agent (e.g., reduction in the average number of daily bowel movements) AND
- 4. The requested agent will be used in combination with a somatostatin analog (e.g., Sandostatin [octreotide], Sandostatin LAR [octreotide], Somatuline Depot [lanreotide])

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Xgeva PA

Drug Name(s)

Xgeva

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of multiple myeloma AND BOTH of the following:
 - i. The requested agent will be used for the prevention of skeletal-related events AND
 - ii. ONE of the following:
 - 1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
 - 2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
 - 3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR
 - B. Patient has a diagnosis of prostate cancer AND ALL of the following:
 - i. The requested agent will be used for the prevention of skeletal-related events AND
 - ii. Patient has bone metastases AND
 - iii. ONE of the following:
 - 1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
 - 2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
 - 3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

- C. Patient has a solid tumor cancer diagnosis (e.g., thyroid, non-small cell lung, kidney cancer, or breast cancer) AND ALL of the following:
 - i. The requested agent will be used for the prevention of skeletal-related events AND
 - ii. Patient has bone metastases AND

- iii. ONE of the following:
 - 1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
 - 2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
 - 3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR
- D. Patient has a diagnosis of giant cell tumor of bone AND ONE of the following:
 - i. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
 - ii. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
 - iii. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR
- E. Patient has a diagnosis of hypercalcemia of malignancy AND
- 2. Patient will NOT be using the requested agent in combination with Prolia (denosumab) AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Xifaxan PA

Drug Name(s)

Xifaxan

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

- 1. Patient has ONE of the following:
 - a. A diagnosis of irritable bowel syndrome with diarrhea (IBS-D) OR
 - b. A diagnosis of hepatic encephalopathy [reduction in risk of overt hepatic encephalopathy (HE) recurrence] OR
 - c. BOTH of the following:
 - i. A diagnosis of traveler's diarrhea (TD) AND
 - ii. The traveler's diarrhea is caused by noninvasive strains of Escherichia coli

Age Restriction:

For diagnosis of traveler's diarrhea (TD), patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Xolair PA

Drug Name(s)

Xolair

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of moderate to severe persistent asthma AND ALL of the following:
 - i. ONE of the following:
 - a. Patient is 6 to less than 12 years of age AND BOTH of the following:
 - I. Patient's pretreatment IgE level is 30 IU/mL to 1300 IU/mL AND
 - II. Patient's weight is 20 kg to 150 kg OR
 - b. Patient is 12 years of age or over AND BOTH of the following:
 - I. Patient's pretreatment IgE level is 30 IU/mL to 700 IU/mL AND
 - II. Patient's weight is 30 kg to 150 kg AND
 - iii. Allergic asthma has been confirmed by a positive skin test or in vitro reactivity test (RAST) to a perennial aeroallergen AND
 - iv. ONE of the following:
 - a. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LRTA, LAMA, theophylline) in combination with the requested agent OR
 - b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an asthma control therapy OR
 - B. Patient has a diagnosis of chronic idiopathic urticaria AND BOTH of the following:
 - i. Patient has had over 6 weeks of hives and itching AND
 - ii. ONE of the following:
 - a. Patient has tried and had an inadequate response to maximum tolerable H1 antihistamine therapy OR
 - b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to H1 antihistamine therapy OR
 - C. Patient has a diagnosis of nasal polyps AND BOTH of the following:
 - i. ONE of the following:
 - a. Patient has tried and had an inadequate response to an intranasal corticosteroid OR
 - b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an intranasal corticosteroid AND
 - ii. ONE of the following:
 - a. The requested agent will be used in combination with an intranasal corticosteroid OR

b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an intranasal corticosteroid OR

Initial criteria continues: see Other Criteria

Age Restriction:

For diagnosis of moderate to severe persistent asthma, patient is 6 years of age or over. For diagnosis of chronic idiopathic urticaria, patient is 12 years of age or over. For diagnosis of nasal polyps, patient is 18 years of age or over.

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, immunologist, otolaryngologist, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal

Other Criteria:

- D. Patient has a diagnosis of IgE-mediated food allergy AND ALL of the following:
 - i. Patient is using the requested agent for the reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods AND
 - ii. IgE-mediated food allergy has been confirmed by an allergy diagnostic test (e.g., skin prick test, serum specific IgE test, oral food challenge) AND
 - iii. Patient will avoid known food allergens while treated with the requested agent AND
 - iv. The requested agent will NOT be used for the emergency treatment of allergic reactions, including anaphylaxis AND
- 2. Patient will NOT be using the requested agent in combination with Dupixent or an injectable Interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of moderate to severe persistent asthma AND ALL of the following:
 - i. Patient's weight is within the FDA indicated range for their age (i.e., 20 kg to 150 kg for patients age 6 to less than 12 years and 30 kg to 150 kg for patients 12 years of age or over) AND
 - ii. Patient has had clinical benefit with the requested agent AND
 - iii. ONE of the following:
 - a. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LRTA, LAMA, theophylline) in combination with the requested agent OR
 - b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an asthma control therapy OR
 - B. Patient has a diagnosis of chronic idiopathic urticaria AND the following:

- a. Patient has had clinical benefit with the requested agent OR
- C. Patient has a diagnosis of nasal polyps AND the following:
 - a. Patient has had clinical benefit with the requested agent OR
- D. Patient has a diagnosis of IgE-mediated food allergy AND ALL of the following:
 - a. Patient is using the requested agent for the reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods AND
 - b. Patient has had clinical benefit with the requested agent AND
 - c. Patient will avoid known food allergens while treated with the requested agent AND
 - d. The requested agent will NOT be used for the emergency treatment of allergic reactions, including anaphylaxis AND
- 3. Patient will NOT be using the requested agent in combination with Dupixent or an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Zepatier PA

Drug Name(s)

Zepatier

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND
- 2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
- 3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
- 4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND
- 5. If genotype 1, the patient's subtype has been identified and provided AND
- 6. If genotype 1a, the prescriber has tested the patient for the presence of virus with NS5A resistance-associated polymorphisms AND
- 7. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR
 - D. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported **Other Criteria**:

Zokinvy PA

Drug Name(s)

Zokinvy

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

- 1. ONE of the following:
 - A. BOTH of the following:
 - i. Patient has a diagnosis of Hutchinson-Gilford progeria syndrome (HGPS) AND
 - ii. Genetic testing has confirmed a pathogenic variant in the LMNA gene that results in production of progerin OR
 - B. Patient has a diagnosis of processing-deficient progeroid laminopathy AND ONE of the following:
 - i. Genetic testing has confirmed heterozygous LMNA mutation with progerin-like protein accumulation OR
 - ii. Genetic testing has confirmed homozygous or compound heterozygous ZMPSTE24 mutations

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has a diagnosis of ONE of the following:
 - A. Hutchinson-Gilford progeria syndrome (HGPS) OR
 - B. Processing-deficient progeroid laminopathies with either: heterozygous LMNA mutation with progerin-like protein accumulation OR homozygous or compound heterozygous ZMPSTE24 mutations AND
- 3. Patient has had clinical benefit with the requested agent

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist, geneticist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months